

# LINEAR EMBEDDINGS OF GRASSMANNIANS AND IND-GRASSMANNIANS

IVAN PENKOV AND VALDEMAR TSANOV

ABSTRACT. By a Grassmannian we understand a usual complex Grassmannian or possibly an orthogonal or symplectic Grassmannian. We classify, with few exceptions, linear embeddings of Grassmannians into larger Grassmannians, where the linearity requirement is the condition that the embedding induces an isomorphism on Picard groups. This classification implies that most linear embeddings of Grassmannians are equivariant.

A linear ind-Grassmannian is the direct limit of a chain of linear embeddings of Grassmannians. We conclude the paper by classifying linear embeddings of linear ind-Grassmannians.

Mathematics Subject Classification 2020: 14E25, 14L30, 14M15.

Keywords: *Grassmannian, isotropic Grassmannian, ind-Grassmannian, linear embedding, equivariant embedding.*

## INTRODUCTION

Projective lines on Grassmannians, and more general linearly embedded projective spaces in Grassmannians, are classical objects in projective geometry. On the other hand, the Plücker embedding realizes a Grassmannian as a projective variety. What these constructions have in common is that the sheaf  $\mathcal{O}(1)$  on the ambient variety restricts to  $\mathcal{O}(1)$  on the subvariety. It makes sense to consider embeddings of arbitrary Grassmannians with this property, and following [9] we call an embedding of complex Grassmannians  $X \xrightarrow{\varphi} Y$  *linear* if  $\mathcal{O}_X(1) \cong \varphi^* \mathcal{O}_Y(1)$ . Here we allow  $X$  or  $Y$ , possibly both  $X$  and  $Y$ , to be isotropic Grassmannians. In [9] the linear embeddings of Grassmannians of the same type, i.e., when both  $X$  and  $Y$  are ordinary Grassmannians, or when both  $X$  and  $Y$  are orthogonal or symplectic Grassmannians, have been classified with some exceptions.

A main application of this classification has been the classification of linear ind-Grassmannians [9]. The latter are defined as direct limits of usual finite-dimensional Grassmannians of the same type under certain linear embeddings called standard extensions. Every linear ind-Grassmannian is a homogeneous ind-space for one of the ind-groups  $SL(\infty) = \varinjlim SL(n)$ ,  $SO(\infty) = \varinjlim SO(n)$ ,  $Sp(\infty) = \varinjlim Sp(n)$ . A famous particular case of a linear ind-Grassmannian is the Sato Grassmannian.

In this paper we classify linear embeddings  $X \xrightarrow{\varphi} Y$  of Grassmannians and ind-Grassmannians, with some exceptions in the case of a spinor (ind-)Grassmannian  $Y$ . It is essential that  $X$  and  $Y$  may have different types.

Here is a brief description of the content. In Section 1 we present some main definitions concerning linear embeddings of Grassmannians, and also recall the families of maximal linearly embedded projective spaces in Grassmannians. Next, in Section 2 we show that using the results of [9] one can classify linear embeddings of Grassmannians of different types, for instance embeddings of ordinary Grassmannians into orthogonal Grassmannians, embeddings of symplectic Grassmannians into ordinary Grassmannians, etc. There are three types of orthogonal Grassmannians which behave in a special way and whose consideration is postponed until Section 3. There we construct some special embeddings which enable us to complete the classification. Section 3 is concluded by a list of maximal embeddings of Grassmannians.

Section 4 is devoted to a study of equivariance properties of the linear embeddings. We show that most embeddings are actually equivariant. Non-equivariant embeddings occur only into isotropic Grassmannians, and their images are contained in projective spaces, quadrics, or Grassmannians of isotropic planes. Finally, in Section 5 we use the classification from Sections 2 and 3 to classify the linear embeddings of linear ind-Grassmannians, with the same few exceptions as in the case of finite-dimensional Grassmannians.

Our arguments make essential use of properties of families of linearly embedded projective spaces in Grassmannians. Such families have been studied in detail by Landsberg and Manivel in [7] from a Lie-theoretic point of view, also in the context of general flag varieties. It is conceivable that a combination of the two approaches could have interesting applications in the study of embeddings of flag varieties.

We dedicate this paper to the memory of our late friend Joseph A. Wolf, with sadness that we can no longer share the joy of mathematics with him.

**Acknowledgement:** Both authors have been supported in part by DFG grant PE 980/9-1. V.T. has also been supported by the Bulgarian Ministry of Education and Science, Scientific Programme “Enhancing the Research Capacity in Mathematical Sciences (PIKOM)”, No. DO1-67/05.05.2022.

## 1. BASIC DEFINITIONS AND PRELIMINARIES

The base field is the field of complex numbers  $\mathbb{C}$ . The notation  $*$  indicates dual vector space or pullback of vector bundles along morphisms of algebraic varieties, depending on the context. Symmetric and exterior powers of a vector space  $V$  are denoted respectively by  $S^n V$  and  $\Lambda^n V$ .

**1.1. Grassmannians.** We denote by  $V_n$  an  $n$ -dimensional complex vector space, by  $\mathbb{P}^n$  a complex projective space of dimension  $n$ , and by  $\mathbb{P}(V_n) \cong \mathbb{P}^{n-1}$  the projective space of 1-dimensional subspaces of  $V_n$ . The Grassmannian  $G(m, V_n)$  of  $m$ -dimensional subspaces of  $V_n$ , for  $1 \leq m \leq n-1$ , is an algebraic variety of dimension  $m(n-m)$ . The Plücker embedding

$$G(m, V_n) = \{U \subset V_n : \dim U = m\} \begin{array}{l} \xrightarrow{\pi} \mathbb{P}(\Lambda^m V_n), \\ U \mapsto \Lambda^m U \end{array}$$

realizes  $G(m, V_n)$  as a projective variety. It is a standard fact that the Picard group of  $G(m, V_n)$  is isomorphic to  $\mathbb{Z}$  and is generated by the class of  $\mathcal{O}_{G(m, V_n)}(1)$ , where  $\mathcal{O}_{G(m, V_n)}(1)$  is the line bundle with fibre  $\Lambda^m U^*$  over the point  $U \in G(m, V_n)$ . This and our statements below concerning the Picard groups are well known consequences of theory of algebraic homogeneous spaces. A good reference for this theory is [5]. We have  $\mathcal{O}_{G(m, V_n)}(1) = \pi^* \mathcal{O}_{\mathbb{P}(\Lambda^m V_n)}(1)$ .

For a fixed symmetric or skew-symmetric nondegenerate bilinear form on  $V_n$ , denoted respectively by  $\kappa$  or  $\omega$ , the maximal isotropic subspaces of  $V_n$  have dimension  $\lfloor n/2 \rfloor$ . The Grassmannian of isotropic subspaces of dimension  $m$  in  $V_n$  is, respectively,

$$\begin{aligned} GO(m, V_n) &:= \{U \subset V_n : \dim U = m, \kappa|_{U \times U} = 0\} \xrightarrow{\tau} G(m, V_n), \quad m < n/2 \\ GS(m, V_n) &:= \{U \subset V_n : \dim U = m, \omega|_{U \times U} = 0\} \xrightarrow{\tau} G(m, V_n), \quad m \leq n/2, \end{aligned}$$

We call  $\tau$  the **tautological embedding**.

Note that all above Grassmannians are connected, with one exception: the variety of  $m$ -dimensional (maximal) isotropic subspaces of a  $2m$ -dimensional orthogonal space. It is well known that this variety has two connected components, denoted  $GO^+(m, V_{2m})$  and  $GO^-(m, V_{2m})$ , which are isomorphic to each other and to  $GO(m-1, V_{2m-1})$ . Two maximal isotropic subspaces of  $V_{2m}$  belong to same connected component if and only if their intersection has even codimension in each of them. This can be checked directly, or we refer the reader to [11] where this fact is discussed in the equivalent language of projective subspaces of quadrics. In what follows, we

simplify the notation to  $GO(m, V_{2m}) = GO^+(m, V_{2m})$ , unless both components are necessary, and we refer to these varieties as **spinor Grassmannians**.

The Picard group of  $GS(m, V_n)$  is freely generated by the class of  $\mathcal{O}_{GS(m, V_n)}(1) := \tau^* \mathcal{O}_{G(m, V_n)}(1)$ . The analogous statement holds for  $GO(m, V_n)$  with three exceptions:  $n = 2m, 2m+1$  or  $2m+2$ . In fact, the first two exceptions “coincide”, as the corresponding to isomorphic spinor Grassmannians. The Picard group here, say for  $n = 2m$ , is freely generated by the class of a square root of  $\tau^* \mathcal{O}_{G(m, V_{2m})}(1)$ , and we denote such a square root by  $\mathcal{O}_{GO(m, V_{2m})}(1)$ . The third exception  $GO(m, V_{2m+2})$  is notably different, as its Picard group is isomorphic to  $\mathbb{Z}^2$ ; in what follows we exclude this variety from our discussion. Again, we refer the reader to [5] for relevant properties of Picard groups. Henceforth, by a **Grassmannian** we mean one of the varieties  $G(m, V_n)$ ,  $GO(m, V_n)$ ,  $GS(m, V_n)$ , except  $GO(m, V_{2m+2})$ .

There are a few isomorphisms between Grassmannians, well known and classified by Onishchik [8, Theorem 7.1]. The infinite series of such isomorphisms are two: the above mentioned isomorphisms of spinor Grassmannians and the isomorphisms of odd-dimensional projective spaces  $\mathbb{P}^{2m-1} = G(1, V_{2m})$  and symplectic Grassmannians  $GS(1, V_{2m})$ . In low dimensions, there is a finite number of coincidences, i.e., isomorphisms:  $\mathbb{P}^1 \cong G(1, V_2) \cong GS(1, V_2) \cong GO(1, V_3) \cong GO(2, V_4)$ ,  $G(2, V_4) \cong GO(1, V_6)$ ,  $GS(2, V_4) \cong GO(1, V_5)$ ,  $\mathbb{P}^3 \cong G(1, V_4) \cong GS(1, V_4) \cong GO(2, V_5) \cong GO(3, V_6)$ ,  $GO(1, V_8) \cong GO(4, V_8) \cong GO(3, V_7)$ .

The Grassmannian  $GO(1, V_n)$  is a quadric of dimension  $n - 2$  in  $\mathbb{P}(V_n)$ , and we shall also use the notation  $Q^{n-2}$  for this variety.

For a Grassmannian  $X$  we set

$$V_X := H^0(X, \mathcal{O}_X(1))^*$$

and denote by  $\pi_X$  the natural embedding  $\pi_X : X \hookrightarrow \mathbb{P}(V_X)$ .

**1.2. Linear embeddings.** Let  $X, Y$  be Grassmannians. We start by recalling the definition of a linear embedding  $X \xrightarrow{\varphi} Y$ .

**Definition 1.1.** *An embedding  $X \xrightarrow{\varphi} Y$  is linear if  $\mathcal{O}_X(1) \cong \varphi^* \mathcal{O}_Y(1)$ .*

Clearly, a composition of two embeddings of Grassmannians is linear if and only if both embeddings are linear. A linear embedding  $X \xrightarrow{\varphi} Y$  is said to **factor through a projective space** if it can be written as a composition of embeddings

$$\varphi : X \xrightarrow{\pi_X} \mathbb{P}(V_X) \xrightarrow{\psi} Y .$$

Here  $\psi$  is necessarily linear. Also note that any linear embedding of Grassmannians  $X \xrightarrow{\varphi} Y$  induces a linear embedding of projective spaces

$$(1) \quad \mathbb{P}(V_X) \xrightarrow{\hat{\varphi}} \mathbb{P}(V_Y) .$$

**Example 1.1.** *The definition of  $\mathcal{O}_X(1)$  for a Grassmannian  $X$  given above implies the following.*

- *The Plücker embedding  $G(m, V_n) \hookrightarrow \mathbb{P}(\Lambda^m V_n)$ , and more generally, the embedding  $\pi_X : X \hookrightarrow \mathbb{P}(V_X)$  for any Grassmannian  $X$  is linear as  $V_X = H^0(X, \mathcal{O}_X(1))^*$ .*
- *The tautological embedding  $GO(m, V_n) \xrightarrow{\tau} G(m, V_n)$  is a linear embedding of Grassmannians if and only if  $n \geq 5$  and  $m < \frac{n}{2} - 1$ .*
- *The tautological embedding  $GS(m, V_n) \xrightarrow{\tau} G(m, V_n)$  is linear.*
- *The Veronese embedding  $\text{Ver}_q : \mathbb{P}(V_n) \hookrightarrow \mathbb{P}(S^q V_n)$ ,  $[v] \mapsto [v^q]$  is not linear for  $q \geq 2$ .*

A **minimal projective embedding** of a Grassmannian  $X$  is a projective embedding  $\pi : X \hookrightarrow \mathbb{P}^k$ , such that there does not exist an embedding of  $X$  into  $\mathbb{P}^l$  for  $l < k$ . The embedding

$\pi_X$  is a minimal projective embedding. Every minimal projective embedding of  $X$  is linear and has the form  $X \xrightarrow{\pi_X} \mathbb{P}(V_X) \xrightarrow{\alpha} \mathbb{P}(V)$  for a suitable isomorphism  $\alpha$ .

A **projective space on  $Y$**  is a linearly embedded  $\mathbb{P}^k \subset Y$  for some  $k \geq 1$ . These are exactly the subvarieties of  $Y$  sent by  $\pi_Y$  to projective subspaces of  $\mathbb{P}(V_Y)$ . An embedding of Grassmannians  $\varphi : X \rightarrow Y$  is linear if and only if it sends any projective space on  $X$  to a projective space on  $Y$ . A **quadric on  $Y$**  is a linearly embedded quadric  $GO(1, V_k) \subset Y$  for some  $k$ .

Next we recall the definitions of various types of linear embeddings.

**Definition 1.2.** ([9]) *An embedding  $G(m, V_n) \xrightarrow{\sigma} G(l, V_s)$  is said to be a **strict standard extension** if  $\sigma$  is given by*

$$(2) \quad \sigma(U) = U \oplus V''$$

for some isomorphism  $V_s \cong V_n \oplus V'$  and a fixed subspace  $V'' \subset V'$  of dimension  $l - m$ .

An embedding  $G(m, V) \xrightarrow{\varphi} G(l, W)$  is a **standard extension** if it fits into a commutative diagram

$$\begin{array}{ccc} G(m, V) & \xrightarrow{\varphi} & G(l, W) \\ i_1 \downarrow & & \downarrow i_2 \\ G(m', V_n) & \xrightarrow{\sigma} & G(l', V_s), \end{array}$$

where  $i_1, i_2$  are isomorphisms and  $\sigma$  is a strict standard extension.

An embedding  $GO(m, V_n) \xrightarrow{\sigma} GO(l, V_s)$  with  $n - 2m \neq 2$ ,  $s - 2l \neq 2$  is a **standard extension** if  $\sigma$  is given by formula (2) for some orthogonal isomorphism  $V_s \cong V_n \oplus V'$  and a fixed isotropic subspace  $V'' \subset V'$ , where in addition we assume  $l = \lfloor \frac{s}{2} \rfloor$  whenever  $m = \lfloor \frac{n}{2} \rfloor$ .

An embedding  $GS(m, V_n) \xrightarrow{\sigma} GS(l, V_s)$  is a **standard extension** if it is given by formula (2) for some symplectic isomorphism  $V_s \cong V_n \oplus V'$  and for a fixed isotropic subspace  $V'' \subset V'$ .

In what follows we call a **standard quadric** on an orthogonal Grassmannian  $Y = GO(l, V_s)$  the image of a standard extension  $\sigma : GO(1, V_n) \hookrightarrow GO(l, V_s)$ . By analogy, a **standard symplectic projective space** on  $GS(l, V_s)$  is the image of a standard extension  $\sigma : GS(1, V_n) \hookrightarrow GS(l, V_s)$ .

**Definition 1.3.** ([9]) *An embedding  $G(l, V_n) \xrightarrow{\iota} GO(l, V_s)$  is an **isotropic extension** if  $l < n \leq \lfloor \frac{s}{2} \rfloor$ , and there exists an isotropic subspace  $W \subset V_s$  and an isomorphism  $f : V_n \cong W$  such that*

$$\iota(U) = f(U) \subset W \quad \text{for } U \in G(l, V_n).$$

An isotropic extension is **minimal** if  $W$  is a maximal isotropic subspace.

Isotropic extensions and minimal isotropic extensions  $G(l, V_n) \xrightarrow{\iota} GS(l, V_s)$  are defined analogously.

**Lemma 1.1.** *Standard and isotropic extensions are linear embeddings.*

*Proof.* The statement is straightforward, but note that the condition on dimensions in the case of orthogonal Grassmannians is essential for linearity.  $\square$

**Definition 1.4.** ([9]) *A **combination of standard and isotropic extensions** is a sequence of embeddings*

$$GO(m, V_n) \xrightarrow{\tau} G(m, V_n) \xrightarrow{\sigma} G(m', V_{n'}) \xrightarrow{\iota} GO(m', V_{n''})$$

or

$$GS(m, V_n) \xrightarrow{\tau} G(m, V_n) \xrightarrow{\sigma} G(m', V_{n'}) \xrightarrow{\iota} GS(m', V_{n''}),$$

where  $\tau$  is a tautological embedding,  $\sigma$  is a standard extension, and  $\iota$  is an isotropic extension.

A mixed combination of standard and isotropic extensions is a sequence of embeddings

$$GO(m, V_n) \xrightarrow{\tau} G(m, V_n) \xrightarrow{\sigma} G(m', V_{n'}) \xrightarrow{\iota} GS(m', V_{n''})$$

or

$$GS(m, V_n) \xrightarrow{\tau} G(m, V_n) \xrightarrow{\sigma} G(m', V_{n'}) \xrightarrow{\iota} GO(m', V_{n''}),$$

where  $\tau$  is a tautological embedding,  $\sigma$  is a standard extension, and  $\iota$  is an isotropic extension.

**Lemma 1.2.** *Let  $X$  be a Grassmannian and  $x, y \in X$  be two points. Then there exist finitely many projective lines  $L_1, \dots, L_k$  on  $X$  such that  $x, y \in L_1 \cup \dots \cup L_k$ .*

*Proof.* Let  $X := G(m, V)$  and let  $U, U'$  be two points of  $X$ . Then  $U$  and  $U'$  can be connected by a sequence of points  $U = U_1, U_2, \dots, U_s = U'$  on  $X$  such that  $\dim(U_i \cap U_{i+1}) = m - 1$  for  $i = 1, \dots, s - 1$ . Since  $U_i$  and  $U_{i+1}$  lie on a projective line on  $X$ , the statement follows. The cases of orthogonal and symplectic Grassmannians are similar.  $\square$

**1.3. Maximal projective spaces on Grassmannians.** As a prerequisite, we need to recall the descriptions and some intersection properties of maximal projective spaces on Grassmannians. An alternative description of the families of (maximal) projective spaces on Grassmannians, and more general flag varieties, can be found in [7].

**1.3.1. Ordinary Grassmannians.** On a projective space there is a single maximal projective space - the space itself. In the Grassmannian  $G(m, V_n)$  with  $1 < m < n - 1$  there are two connected families of maximal projective spaces. A space from the first family is determined by an  $(m + 1)$ -dimensional subspace  $U_{m+1} \subset V_n$ , and has the form

$$\mathbb{P}_{U_{m+1}}^m := \{U \in G(m, V_n) : U \subset U_{m+1}\} \cong \mathbb{P}^m.$$

A space from the second family is determined by an  $(m - 1)$ -dimensional subspace  $U_{m-1} \subset V_n$ , and has the form

$$\mathbb{P}_{U_{m-1}}^{n-m} := \{U \in G(m, V_n) : U_{m-1} \subset U\} \cong \mathbb{P}^{n-m}.$$

The intersection of any two spaces from the same family is empty or equals a point. The intersection of two spaces from different families is empty or is a projective line.

**1.3.2. Symplectic Grassmannians.** On the Grassmannian  $GS(m, V_{2n})$  with  $1 < m < n$  there are two connected families of maximal projective spaces. A space from the first family is determined by an  $(m + 1)$ -dimensional isotropic subspace  $U_{m+1} \subset V_{2n}$ , and has the form

$$\mathbb{P}_{U_{m+1}}^m := \{U \in GS(m, V_{2n}) : U \subset U_{m+1}\} \cong \mathbb{P}^m.$$

A space from the second family is determined by an  $(m - 1)$ -dimensional isotropic subspace  $U_{m-1} \subset V_{2n}$ , and has the form

$$\mathbb{P}_{U_{m-1}}^{2(n-m)-1} := \{U \in GS(m, V_{2n}) : U_{m-1} \subset U\} \cong \mathbb{P}^{2(n-m)-1}.$$

The spaces  $\mathbb{P}_{U_{m-1}}^{2(n-m)-1}$  are exactly the maximal standard symplectic projective spaces on  $GS(m, V_{2n})$ .

The intersection of two maximal projective spaces on  $GS(m, V_{2n})$  belonging to the same family is empty or equals a point. The intersection of two space of different families is empty or is a projective line. Every maximal projective space on the Grassmannian  $GS(m, V_{2m})$  is a standard symplectic projective line determined by an  $(m - 1)$ -dimensional isotropic subspace  $U_{m-1} \subset V_{2m}$ , and has the form

$$\mathbb{P}_{U_{m-1}}^1 := \{U \in GS(m, V_{2n}) : U_{m-1} \subset U\} \cong \mathbb{P}^1.$$

1.3.3. *Quadrics.* All maximal projective spaces on a quadric  $Q^{n-2} = GO(1, V_n) \subset \mathbb{P}(V_n)$  have dimension  $\lfloor n/2 \rfloor - 1$  and are exactly the projectivizations of maximal isotropic subspaces of  $V_n$ .

For  $n = 2r + 1$ , the maximal projective spaces on  $Q^{2r-1}$  are parametrized by the spinor Grassmannian  $GO(r, V_n)$ . The intersection of two distinct maximal projective spaces on  $Q^{2r-1}$  can be empty or be a projective space of any dimension between 0 and  $r - 1$ . For  $n = 2r$ , there are two connected families of maximal projective spaces on  $Q^{n-2}$ , each parametrized by the spinor Grassmannian  $GO(r, V_n)$ . Two different spaces from the same family intersect in a copy of  $\mathbb{P}^{r-1-2k}$  for some  $k \geq 1$ , and an empty intersection occurs if and only if  $r$  is even. Two spaces from different families intersect in a copy of  $\mathbb{P}^{r-2k}$  for some  $k \geq 1$ , and an empty intersection occurs if and only if  $r$  is odd.

1.3.4. *Generic orthogonal Grassmannians.* On the Grassmannian  $GO(m, V_n)$  with  $1 < m < \lfloor n/2 \rfloor - 1$ , there are two types of maximal projective spaces. The first type constitutes a single connected family, each of whose members is determined by an  $(m + 1)$ -dimensional isotropic subspace  $U_{m+1} \subset V_n$  and has the form

$$\mathbb{P}_{U_{m+1}}^m := \{U \in GO(m, V_n) : U \subset U_{m+1}\} \cong \mathbb{P}^m.$$

The intersection of two distinct projective spaces of this type is either empty or is a point.

Maximal projective spaces on  $GO(m, V_n)$  of the second type are maximal projective spaces on maximal standard quadrics on  $GO(m, V_n)$ . These maximal projective spaces form one or two connected families depending on the parity of  $n$ . A maximal standard quadric on  $GO(m, V_n)$  is determined by an  $(m - 1)$ -dimensional isotropic subspace  $U_{m-1} \subset V_n$ , and has the form

$$Q_{U_{m-1}}^{n-2m} := \{U \in GO(m, V_n) : U_{m-1} \subset U\} \cong Q^{n-2m}.$$

The intersection of two distinct maximal standard quadrics on  $GO(m, V_n)$  is either empty or is a point. The intersection of maximal projective spaces on  $GO(m, V_n)$  contained in one maximal standard quadric is as described in Subsection 1.3.3.

The intersection of a maximal projective space  $\mathbb{P}_{U_{m+1}}^m$  and a maximal standard quadric  $Q_{U_{m-1}}^{n-2m-2}$  is empty or equals a point.

**Proposition 1.3.** *Let  $GO(m, V_n)$  be identified with the image of its tautological embedding in  $G(m, V_n)$ , assuming  $m < \frac{n}{2} - 1$ . If  $P$  is a maximal projective space on  $G(m, V_n)$  then there are four possibilities for the intersection  $P \cap GO(m, V_n)$ : it is empty; it is a point; it is a maximal projective space on  $GO(m, V_n)$ ; it is a maximal standard quadric on  $GO(m, V_n)$ .*

1.3.5. *The Grassmannians  $GO(n - 1, V_{2n+1})$ .* The maximal projective spaces on  $X := GO(n - 1, V_{2n+1})$  form a single connected family parametrized by the spinor Grassmannian of maximal isotropic subspaces of  $V_{2n+1}$ :

$$\mathbb{P}_{U_n}^{n-1} := \{U \in X : U \subset U_n\}, \quad U_n \in GO(n, V_{2n+1}).$$

The intersection of two distinct maximal projective spaces on  $X$  is a point or is empty, i.e., for  $U_n \neq U'_n$  we have

$$\mathbb{P}_{U_n}^{n-1} \cap \mathbb{P}_{U'_n}^{n-1} = \begin{cases} \{U_n \cap U'_n\} & \text{if } \dim U_n \cap U'_n = n - 1, \\ \emptyset & \text{if } \dim U_n \cap U'_n < n - 1. \end{cases}$$

The maximal standard quadrics on  $X$  are parametrized by  $GO(n - 2, V_{2n+1})$ , and have the form

$$Q_{U_{n-2}}^3 := \{U \in X : U_{n-2} \subset U\}.$$

The intersection of two distinct maximal standard quadrics on  $X$  is empty or is a point. The intersection of  $Q_{U_{n-2}}^3$  and  $\mathbb{P}_{U_n}^{n-1}$  is empty if  $U_{n-2} \not\subset U_n$ , and equals the projective line

$$\mathbb{P}_{U_{n-2} \subset U_n}^1 := \{U \in X : U_{n-2} \subset U \subset U_n\}$$

if  $U_{n-2} \subset U_n$ . Every projective line on  $X$  is the intersection of a unique maximal standard quadric and a unique maximal projective space.

1.3.6. *Spinor Grassmannians.* Let  $X := GO(m, V_{2m}) = GO^+(m, V_{2m})$  for  $m \geq 5$ . Two distinct points  $V, W \in X$  lie on a projective line on  $X$  if and only if  $\dim V \cap W = m - 2$ . Thus any projective line on  $X$  is determined uniquely by an  $(m - 2)$ -dimensional isotropic subspace, and has the form

$$(3) \quad \mathbb{P}_{U_{m-2}}^1 := \{W \in GO(m, V_{2m}) : U_{m-2} \subset W\} \quad \text{for} \quad U_{m-2} \in GO(m - 2, V_{2m}) .$$

There are two families of maximal projective spaces on  $GO(m, V_{2m})$ : one family of  $\mathbb{P}^{m-1}$ -s and one family of  $\mathbb{P}^3$ -s, parametrized respectively by  $GO^-(m, V_{2m})$  and  $GO(m - 3, V_{2m})$ . A space from the first family has the form

$$\mathbb{P}_{\bar{U}_m}^{m-1} := \{W \in GO(m, V_{2m}) : \dim W \cap \bar{U}_m = m - 1\} \cong \mathbb{P}^{m-1} , \quad \bar{U}_m \in GO^-(m, V_{2m}) .$$

A space from the second family has the form

$$\mathbb{P}_{U_{m-3}}^3 := \{W \in GO(m, V_{2m}) : U_{m-3} \subset W\} \cong \mathbb{P}^3 , \quad U_{m-3} \in GO(m - 3, V_{2m}) .$$

The intersection of two distinct spaces from the family of maximal  $\mathbb{P}^{m-1}$ -s is empty or is a projective line, i.e., for a pair of distinct subspaces  $\bar{U}_m, \bar{U}'_m \in GO^-(m, V_{2m})$  we have

$$\mathbb{P}_{\bar{U}_m}^{m-1} \cap \mathbb{P}_{\bar{U}'_m}^{m-1} = \begin{cases} \emptyset & \text{if } \dim \bar{U}_m \cap \bar{U}'_m < m - 2 , \\ \mathbb{P}_{\bar{U}_m \cap \bar{U}'_m}^1 & \text{if } \dim \bar{U}_m \cap \bar{U}'_m = m - 2 . \end{cases}$$

There are three possibilities for the intersection of two distinct spaces from the family of maximal  $\mathbb{P}^3$ -s: it is empty, it is a point, or it is a projective line. Indeed,  $\mathbb{P}_{U_{m-3}}^3$  and  $\mathbb{P}_{U'_{m-3}}^3$  intersect if and only if the sum  $U := U_{m-3} + U'_{m-3}$  is contained in  $U_m$  for some  $U_m \in X$ . If  $U_{m-3} \subset U_m$  for some  $U_m \in X$  and  $U_{m-3} \neq U'_{m-3}$ , then the intersection  $U \cap U_m$  has codimension 0, 1 or 2 in  $U_m$ . Hence

$$\mathbb{P}_{U_{m-3}}^3 \cap \mathbb{P}_{U'_{m-3}}^3 = \{W \in GO(m, V_{2m}) : W \supset U\} = \begin{cases} \emptyset & \text{if } U \not\subset W \forall W \in X , \\ \{U_m\} & \text{if } \dim U \geq m - 1 , U \subset U_m \in X , \\ \mathbb{P}_U^1 & \text{if } \dim U = m - 2 . \end{cases}$$

There are three possibilities for the intersection of  $\mathbb{P}_{\bar{U}_m}^{m-1}$  and  $\mathbb{P}_{U_{m-3}}^3$ : it is empty, a point, or a projective plane. More precisely,

$$\mathbb{P}_{\bar{U}_m}^{m-1} \cap \mathbb{P}_{U_{m-3}}^3 = \begin{cases} \emptyset & \text{if } \dim(U_{m-3} + \bar{U}_m) > m + 1 , \\ \{V_m\} & \text{if } \begin{cases} \dim(U_{m-3} + \bar{U}_m) = m + 1 , \\ U_{m-3} + \bar{U}_m = V_m + \bar{U}_m \text{ for some } V_m \in X , \end{cases} \\ \mathbb{P}_{U_{m-3} \subset \bar{U}_m}^2 & \text{if } U_{m-3} \subset \bar{U}_m , \end{cases}$$

where

$$\mathbb{P}_{U_{m-3} \subset \bar{U}_m}^2 := \{W \in GO(m, V_{2m}) : U_{m-3} \subsetneq W \cap \bar{U}_m\} \cong \mathbb{P}^2 .$$

For every  $\mathbb{P}^2$  on  $GO(m, V_{2m})$  containing a point  $V_m$ , there exists a unique flag  $U_{m-3} \subset \bar{U}_m$  such that  $\dim V_m \cap \bar{U}_m = m - 1$  and  $\mathbb{P}^2 = \mathbb{P}_{U_{m-3} \subset \bar{U}_m}^2$ .

## 2. CLASSIFICATIONS OF LINEAR EMBEDDINGS OF GRASSMANNIANS

2.1. **Linear embeddings into quadrics.** Let  $X \subset \mathbb{P}(V_X)$  be a Grassmannian identified with its image under the minimal projective embedding  $\pi_X$ . It is well known that the ideal  $I(X)$  in the homogeneous coordinate ring  $\mathbb{C}[V_X]$  is generated by its degree-two component  $I_2(X)$ , which is in turn spanned by (in most cases degenerate) quadratic polynomials, see e.g. [6, Theorem 16.2.2.6]. For  $X = G(m, V_n)$  these quadratic polynomials are the original Plücker relations. The ideal of  $X$  is of course trivial if and only if  $X = \mathbb{P}(V_X)$ .

**Proposition 2.1.** *Let  $X \subset \mathbb{P}(V_X)$  be a projectively embedded Grassmannian. For every nonzero  $p \in I_2(X)$  there exists a linear embedding  $X \xrightarrow{\kappa_p} Q = GO(1, V_n)$  with  $n = 2 \dim V_X - \text{rank } p$ , which does not factor through a projective space or through a quadric of smaller dimension. Furthermore, if  $Q \subset \mathbb{P}(V_n)$  is tautologically embedded and  $q \in I_2(Q)$  is a generator of the ideal of  $Q$ , then  $\kappa_p$  extends to a linear embedding  $\tilde{\kappa}_p : \mathbb{P}(V_X) \hookrightarrow \mathbb{P}(V_n)$  and  $\tilde{\kappa}_p^* q = ap$  for some nonzero scalar  $a$ .*

*Conversely, every linear embedding  $X \xrightarrow{\varphi} Q^{s-2} = GO(1, V_s)$ , which does not factor through a projective space, factors as a composition  $\varphi : X \xrightarrow{\kappa_p} Q \xrightarrow{\sigma} Q^{s-2}$  for some nonzero  $p \in I_2(X)$  and some standard extension of quadrics  $\sigma$ .*

*Proof.* Let  $p \in I_2(X)$  be a nonzero element and  $r$  be its rank. Then  $r \geq 2$  since  $X$  is not contained in a hyperplane on  $\mathbb{P}(V_X)$ . The space  $V_X$  admits a decomposition  $V_X = V_r \oplus U$  such that the restriction of  $p$  to  $V_r$  is non-degenerate and  $U$  is an isotropic space for  $p$ . Set  $V_n := V_X \oplus U^* = V_r \oplus U \oplus U^*$ . Pick a non-degenerate quadratic polynomial  $q \in \mathbb{C}[V_n]_2$  whose restriction to  $V_X$  equals  $p$  and which vanishes on  $U^*$ . The quadric  $Q \subset \mathbb{P}(V_n)$  defined by the vanishing of  $q$  contains  $X$ . Furthermore,  $X$  is not contained in a projective space on  $Q$  because  $\mathbb{P}(V_X)$  is the minimal projective subspace of  $\mathbb{P}(V)$  containing  $X$  and  $q$  does not vanish on  $\mathbb{P}(V_X)$ . To show that  $X$  is not contained in a smaller quadric  $\tilde{Q} \subset Q$ , note that  $\mathbb{P}(V_X) \subset \mathbb{P}(V_{\tilde{Q}}) \subset \mathbb{P}(V_n)$  whenever  $X \subset \tilde{Q} \subset Q$ . On the other hand,  $V_X$  contains the orthogonal space  $W^\perp$  to a maximal  $q$ -isotropic subspace  $W \subset V_n$ , and  $W^\perp$  is not contained in a proper subspace of  $V_n$  on which the restriction of  $q$  is non-degenerate. Hence,  $X \subset \tilde{Q} \subset Q$  implies  $\tilde{Q} = Q$ . This proves the first part of the proposition.

For the converse statement, consider a linear embedding  $X \xrightarrow{\varphi} Q^{s-2}$  which does not factor through a projective space. Let  $Q^{n-2} \subset Q^{s-2}$  be a minimal quadric containing  $\varphi(X)$  and embedded in  $Q^{s-2}$  by a standard extension. Let  $\mathbb{P}(V_X) \xrightarrow{\hat{\varphi}} \mathbb{P}(V_n) \subset \mathbb{P}(V_s)$  be the corresponding linear embeddings of projective spaces. Then  $V_n$  is a minimal non-degenerate subspace of  $V_s$  containing  $V_X$ . Denote by  $q \in I_2(Q^{n-2})$  a generator of the ideal of  $Q^{n-2}$  in  $\mathbb{C}[V_n]$  and let  $p := \hat{\varphi}^* q$ . The element  $p \in I_2(X)$  is nonzero because  $\varphi$  does not factor through a projective space, the equality  $n = 2 \dim V_X - \text{rank } p$  holds, and the embedding of  $X$  into  $Q^{n-2}$  is the embedding  $\kappa_p$ . This completes the proof.  $\square$

The above proposition and its proof have the following two immediate corollaries.

**Corollary 2.2.** *Let  $X \xrightarrow{\varphi} Y$  be a linear embedding of Grassmannians and  $\mathbb{P}(V_X) \xrightarrow{\hat{\varphi}} \mathbb{P}(V_Y)$  be the induced linear embedding of projective spaces. Let  $p_1 \in I_2(X)$ ,  $p_2 \in I_2(Y)$  be nonzero elements and  $\kappa_{p_1}^X, \kappa_{p_2}^Y$  be the respective embeddings of  $X$  and  $Y$  into quadrics. Then the following statements are equivalent:*

- (i) *The embedding  $\kappa_{p_1}^X$  factors as  $\kappa_{p_1}^X = \kappa_{p_2}^Y \circ \varphi$ .*
- (ii) *The equalities  $\hat{\varphi}^* p_2 = ap_1$  (for some nonzero scalar  $a$ ) and  $\dim V_X - \text{rank } p_1 = 2 \dim V_Y - \text{rank } p_2$  hold.*

**Corollary 2.3.** *For any Grassmannian  $X$  let  $r_X := \max\{\text{rank } p : p \in I_2(X)\}$ . Then the minimal number  $n$  for which a linear embedding  $X \hookrightarrow Q^{n-2}$  exists is equal to  $2 \dim V_X - r_X$ . The minimal projective embedding  $X \hookrightarrow \mathbb{P}(V_X)$  factors through a quadric if and only if  $r_X = \dim V_X$ .*

**2.2. Linear embeddings of Grassmannians: generic case.** We now proceed with the main steps of our classification of linear embeddings of Grassmannians. First we recall the following theorem.

**Theorem 2.4.** ([9, Theorem 3.1], Linear embeddings of Grassmannians having the same type)

- (i) *Every linear embedding  $G(m, V_n) \hookrightarrow G(l, V_s)$  is either a standard extension or factors through a projective space.*

- (ii.1) For a linear embedding  $\varphi : GO(m, V_n) \hookrightarrow GO(l, V_s)$ , where either  $l \leq \frac{s}{2} - 2$  and  $m < \frac{n}{2} - 2$ , or both  $n, s$  are odd and  $0 < \lfloor \frac{s}{2} \rfloor - l \leq \lfloor \frac{n}{2} \rfloor - m \leq 2$ , there are four options:  $\varphi$  is a standard extension;  $\varphi$  is a combination of standard and isotropic extensions;  $\varphi$  factors through a projective space;  $\varphi$  factors through a standard quadric but not through a projective space.
- (ii.2) For a linear embedding  $\varphi : GO(\lfloor \frac{n}{2} \rfloor, V_n) \hookrightarrow GO(\lfloor \frac{s}{2} \rfloor, V_s)$  there are two options:  $\varphi$  is a standard extension;  $\varphi$  factors through a projective space.
- (iii) For a linear embedding  $\varphi : GS(m, V_n) \hookrightarrow GS(l, V_s)$  there are three options:  $\varphi$  is a standard extension;  $\varphi$  factors through a projective space;  $\varphi$  is a combination of standard and isotropic extensions.

*Proof.* The statement given here covers a few cases which are omitted in Theorem 3.1 in [9], namely in part (ii) of this theorem the hypothesis is  $l < \lfloor \frac{s}{2} \rfloor - 2$  and  $m < \lfloor \frac{n}{2} \rfloor - 2$  instead of our hypothesis  $l \leq \frac{s}{2} - 2$  and  $m < \frac{n}{2} - 2$ . The proof of Theorem 3.1 in [9] is however valid in the additional cases as well.  $\square$

Our main result in this subsection is the following addition to Theorem 2.4.

**Theorem 2.5.** (Linear embeddings of Grassmannians having different types)

- (i) A linear embedding  $G(m, V_n) \hookrightarrow GO(l, V_s)$  with  $s \neq 2l, 2l + 1, 2l + 2$ , which does not factor through a projective space or a standard quadric, is a composition

$$G(m, V_n) \xrightarrow{\sigma} G(l, V_{n'}) \xrightarrow{\iota} GO(l, V_s) ,$$

where  $\sigma$  is a standard extension and  $\iota$  is an isotropic extension.

- (ii) A linear embedding  $G(m, V_n) \hookrightarrow GS(l, V_s)$  is a composition

$$G(m, V_n) \xrightarrow{\sigma} G(l, V_{n'}) \xrightarrow{\iota} GS(l, V_s) ,$$

where  $\sigma$  is a standard extension and  $\iota$  is an isotropic extension, or factors through a projective space.

- (iii) A linear embedding  $GO(m, V_n) \hookrightarrow G(l, V_s)$  with  $n \geq 7$  and  $m < \lfloor \frac{n}{2} \rfloor - 2$ , which does not factor through a projective space, is a composition

$$GO(m, V_n) \xrightarrow{\tau} G(m, V_n) \xrightarrow{\sigma} G(l, V_s) ,$$

where  $\tau$  is a tautological embedding and  $\sigma$  is a standard extension.

- (iv) A linear embedding  $GS(m, V_n) \hookrightarrow G(l, V_s)$ , which does not factor through a projective space, is a composition

$$GS(m, V_n) \xrightarrow{\tau} G(m, V_n) \xrightarrow{\sigma} G(l, V_s) ,$$

where  $\tau$  is a tautological embedding and  $\sigma$  is a standard extension.

- (v) A linear embedding  $GO(m, V_n) \hookrightarrow GS(l, V_s)$ , where  $n \geq 7$  and  $m < \lfloor \frac{n}{2} \rfloor - 2$ , is a mixed combination of standard and isotropic extensions, or factors through a projective space. The same holds for a linear embedding  $GS(m, V_n) \hookrightarrow GO(l, V_s)$ , where  $7 \leq s \neq 2l, 2l + 1, 2l + 2$ , with the additional possibility for it to factor through a standard quadric.

*Proof.* For part (i) we may assume that  $m \neq 1, n - 1$  since the statement is already known for embeddings of projective spaces, see Subsection 1.3.4. We consider the composition

$$G(m, V_n) \xrightarrow{\varphi} GO(l, V_s) \xrightarrow{\tau} G(l, V_s) ,$$

where  $\varphi$  is a given linear embedding and  $\tau$  is the tautological embedding. By Theorem 2.4,(i) the embedding  $\tau \circ \varphi$  is either a standard extension or factors through a projective space on  $G(l, V_s)$ . Let us first assume that  $\tau \circ \varphi$  is a strict standard extension. Then the image of  $\tau \circ \varphi$  consists of all  $l$ -dimensional subspaces of  $V_s = V_n \oplus V$  of the form  $U_m \oplus W$  for some fixed  $(l - m)$ -dimensional subspace  $W \subset V$  and  $U_m \in G(m, V_n)$ . By hypothesis the image of  $\tau \circ \varphi$

lies in  $GO(l, V_s)$ , so each subspace  $U_m \oplus W$  is isotropic in  $V_s$ . Therefore the entire space  $V_n \oplus W$  is isotropic in  $V_s$ , and hence  $\varphi$  is the composition

$$G(m, V_n) \xrightarrow{\varphi'} G(l, V_n \oplus W) \xrightarrow{\iota} GO(l, V_s)$$

where  $\varphi'(U_m) = U_m \oplus W$  and  $\iota(U_m \oplus W) = U_m \oplus W \in GO(l, V_s)$ . The claim follows in this case.

We now suppose that  $\tau \circ \varphi$  is a non-strict standard extension. Then the composition

$$G(n - m, V_n^*) \xrightarrow{\delta} G(m, V_n) \xrightarrow{\varphi} GO(l, V_s) \xrightarrow{\tau} G(l, V_s)$$

of  $\tau \circ \varphi$  with the duality isomorphism  $\delta$  is a strict standard extension. Our claim is already proven for strict standard extensions. Thus  $\varphi \circ \delta = \iota \circ \sigma'$  where  $\sigma' : G(n - m, V_n^*) \rightarrow G(m', V_{n'})$  is a standard extension and  $\iota : G(m', V_{n'}) \rightarrow GO(l, V_s)$  is an isotropic extension. Then  $\sigma := \sigma' \circ \delta^{-1} : G(m, V_n) \rightarrow G(m', V_{n'})$  is a standard extension and  $\varphi = \iota \circ \sigma$  as required.

It remains to consider the case where  $\tau \circ \varphi$  factors through a projective space, i.e.,  $\tau \circ \varphi = \lambda \circ \psi$  where  $G(m, V_n) \xrightarrow{\psi} \mathbb{P}^k \xrightarrow{\lambda} G(l, V_s)$ . Here  $\varphi(G(m, V_n)) \subset \lambda(\mathbb{P}^k) \cap \tau(GO(l, V_s))$ . We can assume that  $\lambda(\mathbb{P}^k)$  is a maximal projective space on  $G(l, V_s)$ , in which case there are two possibilities: either  $k = l$  and  $\lambda(\mathbb{P}^k) = \{U_l \in G(l, V_s) : U_l \subset W\} =: \mathbb{P}_W^l$  for a fixed subspace  $W \subset V_s$  of dimension  $l + 1$ , or  $k = s - l$  and  $\lambda(\mathbb{P}^k) = \{U_l \in G(l, V_s) : U \subset U_l\} =: \mathbb{P}_U^{s-l}$  for a fixed subspace  $U \subset V_s$  of dimension  $l - 1$ .

We claim that in the former case  $W$  is necessarily isotropic. This follows from the observation that  $W$  contains all  $l$ -dimensional spaces  $U_l$  in the image of the embedding  $\tau \circ \varphi$ , hence at least two different isotropic  $l$ -dimensional subspaces. Thus  $\lambda(\mathbb{P}^k) \subset \tau(GO(l, V_s))$  and  $\varphi$  factors through a projective space, which contradicts our assumption.

In the case where  $\lambda(\mathbb{P}^k) = \mathbb{P}_U^{s-l}$  with  $\dim U = l - 1$ , we observe that  $\lambda(\mathbb{P}^k) \cap \tau(GO(l, V_s)) =: Q_U^{s-2l-2}$  is a maximal standard quadric on  $GO(l, V_s)$ , therefore  $\varphi$  factors through this quadric. Since the dimension of  $G(m, V_n)$  is at least 4 we have  $s - 2l \geq 6$ , and hence the quadric  $Q_U^{s-2l-2}$  is a Grassmannian with Picard group isomorphic to  $\mathbb{Z}$  and generated by the restriction of  $\mathcal{O}_{\mathbb{P}_U^{s-l}}(1)$ . By assumption  $\varphi$  does not factor through a projective space on  $GO(l, V_s)$ , so the embedding of  $G(m, V_n)$  in  $Q_U^{s-2l-2}$  is one of the embeddings described in Proposition 2.1. Part (i) is proved.

Part (ii) is analogous to part (i). Here  $\lambda(\mathbb{P}^k) \cap \tau(GS(l, V_s))$  is a projective space and the possibility of factoring through a quadric does not occur.

Part (iii). Let  $GO(m, V_n) \xrightarrow{\varphi} G(l, V_s)$  be a linear embedding. We assume that  $l \leq s/2$ , otherwise the argument below can be applied to the composition of  $\varphi$  with the duality isomorphism  $G(l, V_s) \cong G(s - l, V_s^*)$ . Now the inequality  $n \geq 7$  implies  $l < s - 2$ . By hypothesis we also have  $m < \lfloor \frac{n}{2} \rfloor - 2$ , and hence Theorem 2.4 can be applied to the composition

$$GO(m, V_n) \xrightarrow{\varphi} G(l, V_s) \xrightarrow{\iota_0} GO(l, V_{2s}) ,$$

where  $\iota_0$  is a minimal isotropic extension where  $V_s$  is identified with a maximal isotropic subspace of  $V_{2s}$ . Since the embedding  $\iota_0 \circ \varphi$  factors through the Grassmannian  $G(l, V_s)$ , it cannot be a standard extension, and by Theorem 2.4 there are three options:  $\iota_0 \circ \varphi$  factors through a projective space, factors through a quadric, or is a combination of standard and isotropic extensions of the form

$$GO(m, V_n) \xrightarrow{\tau} G(m, V_n) \xrightarrow{\sigma} G(l, V_r) \xrightarrow{\iota} GO(l, V_{2s}) .$$

If  $\iota_0 \circ \varphi$  factors through a projective space or a standard quadric  $Z \subset GO(l, V_{2s})$ , then the intersection  $Z \cap \iota_0(G(l, V_s))$  is a projective space through which  $\varphi$  factors.

If the third option holds and  $\iota_0 \circ \varphi = \iota \circ \sigma \circ \tau$  then, from the definitions of standard and isotropic extensions, we obtain injective linear maps  $V_n \hookrightarrow V_r \hookrightarrow V_{2s}$  with  $V_r \hookrightarrow V_{2s}$  isotropic. By altering the map  $V_r \hookrightarrow V_{2s}$  without changing the image of  $V_n$  in  $V_{2s}$ , if necessary we can

assume that the image of  $V_r$  is contained in the maximal isotropic subspace  $V_s$  of  $V_{2s}$ . This yields an embedding  $\iota(G(l, V_r)) \xrightarrow{\sigma_1} \iota_0(G(l, V_s))$  which must be a standard extension due to Theorem 2.4,(i), as under our current hypothesis  $\varphi$  does not factor through a projective space. Then the composition  $\sigma_2 := \sigma_1 \circ \sigma : G(m, V_n) \hookrightarrow G(l, V_s)$  is a standard extension satisfying  $\varphi = \sigma_2 \circ \tau$  as required.

Part (iv) is analogous to part (iii).

Part (v). First we consider a linear embedding  $GO(m, V_n) \xrightarrow{\varphi} GS(l, V_s)$ . Let us form the composition

$$GO(m, V_n) \xrightarrow{\varphi} GS(l, V_s) \xrightarrow{\tau} G(l, V_s),$$

where  $\tau$  is the tautological embedding. By (iii),  $\tau \circ \varphi$  either factors through a projective space or can be written as  $\sigma \circ \tau'$ , where

$$GO(m, V_n) \xrightarrow{\tau'} G(m, V_n) \xrightarrow{\sigma} G(l, V_s)$$

with  $\tau'$  tautological and  $\sigma$  a standard extension. In the latter case, the image of  $\sigma \circ \tau'$  lies in  $GS(l, V_s)$  if  $V_s \oplus W$  is isotropic for the symplectic form chosen on  $V_s$ , and we can apply the same argument as in (i) to prove the claim. If  $\tau \circ \varphi$  factors through a projective space  $\lambda(\mathbb{P}^l) \subset G(l, V_s)$ , the image of  $\tau \circ \varphi$  is contained in the intersection  $\lambda(\mathbb{P}^l) \cap \tau(GS(l, V_s))$ , which is in turn a projective space on  $GS(l, V_s)$ . Thus  $\varphi$  factors through a projective space.

The case of a linear embedding  $GS(m, V_n) \xrightarrow{\varphi} GO(l, V_s)$  is analogous, except for the situation where  $\tau \circ \varphi : GS(m, V_n) \rightarrow G(l, V_s)$  factors through a maximal projective space  $\lambda(\mathbb{P}^l) \subset G(l, V_s)$ . Here, as in part (i), the intersection  $\lambda(\mathbb{P}^l) \cap \tau(GO(l, V_s))$  is either a projective space or a standard quadric on  $GO(l, V_s)$ . In the former case  $\varphi$  factors through a projective space. In the latter case, if  $Q \subset GO(l, V_s)$  is a standard quadric containing the image of  $\varphi$ , then the resulting embedding  $\varphi_1 : GS(m, V_n) \hookrightarrow Q$  either factors through a projective space (on  $Q$  and hence on  $GO(l, V_s)$ ) or is one of the embeddings described in Proposition 2.1.  $\square$

**Corollary 2.6.** *If  $X$  is a Grassmannian and  $X \xrightarrow{\varphi} GS(l, V_{2l})$  is a linear embedding then  $X$  is isomorphic to  $GS(m, V_{2m})$  with  $1 \leq m \leq l$ , and the embedding  $\varphi$  is a standard extension.*

*Proof.* The statement follows from the observation that the Grassmannians  $GS(l, V_{2l})$  with  $l \geq 1$  are characterized among all Grassmannians by the property that the maximal projective spaces on them are projective lines, and from Theorem 2.4.  $\square$

**Corollary 2.7.** *Let  $X$  be a Grassmannian. If  $X \xrightarrow{\varphi} GO(l-1, V_{2l+1})$  is a linear embedding which does not factor through a projective space, then  $\varphi$  is a standard extension  $X \cong GO(k-1, V_{2k+1}) \hookrightarrow GO(l-1, V_{2l+1})$  for some  $k \leq l$ .*

*Proof.* Set  $Y = GO(l-1, V_{2l+1})$ . As recalled in Subsection 1.3.5, the maximal projective spaces on  $Y$  form a single family  $\mathbb{P}_{X, U_l}^{l-1}$  parametrized by the points  $U_l$  on spinor Grassmannian  $GO(l, V_{2l+1})$ . The intersection of two distinct maximal projective spaces on  $Y$  is either empty or a point. The only Grassmannians with the latter property are  $GO(k-1, V_{2k+1})$  and  $GS(k, V_{2k})$  for  $k \geq 1$ . This implies that if  $X$  is a Grassmannian not isomorphic to  $GO(k-1, V_{2k+1})$  or  $GS(k, V_{2k})$ , then any linear embedding  $\varphi : X \hookrightarrow Y$  factors through a projective space. Note that the maximal projective spaces on  $Y$  are exactly the images of isotropic extensions  $\iota : G(l-1, U_l) \hookrightarrow Y$  for maximal isotropic subspaces  $U_l \subset V_{2l+1}$ . Since the Grassmannian  $G(l-1, U_l)$  is a projective space, Theorem 2.5,(v) implies that any linear embedding  $GS(k, V_{2k}) \hookrightarrow Y$  factors through a projective space. The case  $X = GO(k-1, V_{2k+1})$  is handled in Theorem 2.4, (ii), and we observe that, unless the embedding  $\varphi$  is a standard extension, it factors through an isotropic extension into  $Y$  and hence through a projective space.  $\square$

**Corollary 2.8.** *Let  $X$  be a non-spinor Grassmannian and  $X \xrightarrow{\varphi} GO(l-2, V_{2l})$  be a linear embedding which does not factor through an isotropic extension  $G(l-2, U_l) \hookrightarrow GO(l-2, V_{2l})$ .*

Then  $X$  is isomorphic to  $GO(k-2, V_{2k})$  or  $GO(k-2, V_{2k-1})$  for some  $k \leq l$ , and  $\varphi$  is a standard extension.

*Proof.* Let  $Y := GO(l-2, V_{2l})$ . We observe that any projective space on  $Y$  is contained in the image of some isotropic extension  $G(l-2, U_l) \hookrightarrow Y$ . Thus the hypothesis implies that  $\varphi$  does not factor through a projective space. A maximal standard quadric on  $Y$  is 4-dimensional, hence if  $\varphi$  factors through a standard quadric then  $X$  is isomorphic to  $\mathbb{P}^1$ ,  $\mathbb{P}^2$ ,  $Q^3$ , or  $Q^4$ . The first two of these cases are excluded by the hypothesis, while in the latter two  $\varphi$  is a standard extension. Further,  $X$  is not an ordinary or symplectic Grassmannian because by Theorem 2.5 every linear embedding of such Grassmannians into  $Y$  factors through an isotropic extension or through a standard quadric. Hence  $X$  is an orthogonal Grassmannian.

The case  $X = GO(m, V_n)$  with  $m < \frac{n}{2} - 2$  is impossible. This follows from Theorem 2.4. Indeed, since there are no standard extensions  $GO(m, V_n) \hookrightarrow GO(l-2, V_{2l})$  with  $m < \frac{n}{2} - 2$ , the only remaining option for  $\varphi$  is to be a combination of standard and isotropic extensions. A contradiction with our assumption.

The case  $X = GO(k-2, V_{2k})$  is considered in [9, Proposition 3.15]. The options for  $\varphi$  can be reduced to the following two:  $\varphi$  is a standard extension or  $\varphi$  factors through an isotropic extension. The proof of [9, Proposition 3.15] extends without substantial alterations to the case of  $X = GO(k-2, V_{2k-1})$ . Indeed, the key step in that proof is to observe that the restriction of  $\varphi$  to a maximal standard quadric  $Q^3 \subset X$  is a standard extension or factors through a projective space. In the latter case  $\varphi$  factors through an isotropic extension, and in the former case  $\varphi$  is a standard extension.  $\square$

**Corollary 2.9.** *Let  $n \geq 7$  and  $Y$  be a non-spinor Grassmannian. Every linear embedding  $GO(1, V_n) \xrightarrow{\varphi} Y$  is a standard extension*

$$GO(1, V_n) \hookrightarrow GO(l, V_s)$$

for  $s - 2l \geq n$ , or factors through a projective space.

*Proof.* The statement follows by examination of the cases occurring in Theorems 2.4, 2.5, and Corollaries 2.7, 2.8.  $\square$

**Remark 2.1.** *Due to the isomorphism  $GO(1, V_5) \cong GS(2, V_4)$ , the three-dimensional quadric admits standard extensions to both orthogonal and symplectic Grassmannians. These exhaust its linear embeddings which do not factor through a projective space. The four-dimensional quadric  $GO(1, V_6)$  is isomorphic to  $G(2, V_4)$ , and hence, besides the standard extensions to orthogonal Grassmannians, it admits standard extensions to ordinary Grassmannians, as well as isotropic extensions to orthogonal and symplectic Grassmannians.*

**2.3. Pullbacks of tautological bundles.** If  $X = G(m, V_n)$ , we denote by  $\mathcal{S}$  (or  $\mathcal{S}_X$  if necessary) the tautological bundle of rank  $m$  on  $X$ . By definition  $\mathcal{S}$  is a subbundle of  $V_n \otimes \mathcal{O}_X$ . Also, we let  $\mathcal{S}^\perp$  denote the bundle  $((V_n \otimes \mathcal{O}_X)/\mathcal{S})^*$  on  $X$ , which is the tautological bundle on  $G(n-m, V_n^*)$ . Here  $(\cdot)^*$  stands for dual bundle. Note that  $(\mathcal{S}^\perp)^\perp = \mathcal{S}$  holds. Below we refer to both  $\mathcal{S}$  and  $\mathcal{S}^\perp$  as **tautological bundles** on  $X$ .

Recall that the Grassmannian  $G(m, V_n)$  represents the functor

$$\mathbb{G}(m, V_n) : \text{Algebraic varieties} \rightarrow \text{Sets}$$

which sends an algebraic variety  $Z$  to the set  $\mathbb{G}(m, V_n)(Z)$  of rank- $m$  subbundles of  $V_n \otimes \mathcal{O}_Z$ . In other words, we have a bijection  $\text{Hom}(Z, G(m, V_n)) \cong \mathbb{G}(m, V_n)(Z)$  via which  $\varphi \in \text{Hom}(Z, G(m, V_n))$  is identified with the subbundle  $\varphi^*\mathcal{S} \subset V_n \otimes \mathcal{O}_Z$ . As a consequence, the subbundle  $\varphi^*\mathcal{S} \subset V_n \otimes \mathcal{O}_Z$  determines the morphism  $\varphi$ . A similar statement holds for the subbundle  $\varphi^*(\mathcal{S}^\perp) \subset V_n^* \otimes \mathcal{O}_Z$ .

If  $X = GO(m, V_n)$  (respectively,  $X = GS(m, V_n)$ ), then  $\mathcal{S}$  stands for the tautological bundle of rank  $m$  on  $X$ . In this situation the bundle  $\mathcal{S}^\perp$  is defined as the subbundle of  $V_n \otimes \mathcal{O}_X$

orthogonal to  $\mathcal{S}$  and we have  $\mathcal{S} \subset \mathcal{S}^\perp \subset V_n \otimes \mathcal{O}_X$ . We reserve the name tautological bundle for  $\mathcal{S}$  but not for  $\mathcal{S}^\perp$ . The Grassmannian  $X$  represents the functor  $\mathbb{G}\mathbb{O}(m, V_n)$  (respectively,  $\mathbb{G}\mathbb{S}(m, V_n)$ ) sending an algebraic variety  $Z$  to the set of rank- $m$  isotropic subbundles of  $V_n \otimes \mathcal{O}_Z$ . Any such subbundle determines a morphism  $\varphi \in \text{Hom}(Z, X)$ .

In the following statement we describe the pullbacks  $\varphi^*\mathcal{S}_Y$  of tautological bundles on  $Y$  for linear embeddings  $X \xrightarrow{\varphi} Y$  of Grassmannians as in Theorems 2.4 and 2.5.

**Proposition 2.10.** *Let  $X \xrightarrow{\varphi} Y$  be a linear embedding, where  $X$  and  $Y$  are non-spinor Grassmannians and in addition  $X$  is not isomorphic to  $GO(m, V_{2m+3})$  or  $GO(m, V_{2m+4})$  for any  $m \geq 3$ .*

- (i) *If the embedding  $\varphi$  does not factor through a projective space or a standard quadric, and  $\mathcal{S}_Y$  is a tautological bundle on  $Y$ , then the pullback  $\varphi^*\mathcal{S}_Y$  is isomorphic to a direct sum of a trivial bundle with  $\mathcal{S}_X$  or  $\mathcal{S}_X^\perp$ , where  $\mathcal{S}_X$  is a tautological bundle on  $X$ .*
- (ii) *If  $Y$  is an ordinary Grassmannian and  $\varphi$  factors through a projective space, then one of pullbacks  $\varphi^*\mathcal{S}_Y$  or  $\varphi^*\mathcal{S}_Y^\perp$  is isomorphic to the direct sum of  $\mathcal{O}_X(-1)$  with a trivial bundle. The same holds for the pullback  $\varphi^*\mathcal{S}_Y$  if  $Y$  is an orthogonal or symplectic Grassmannian, under the assumption that  $\varphi$  factors respectively through a standard quadric or through a standard symplectic projective space.*
- (iii) *If  $Y$  is an orthogonal or symplectic Grassmannian and  $\varphi$  factors through a projective space which is not contained in a standard quadric or, respectively, in a standard symplectic projective space, then  $\varphi^*\mathcal{S}_Y$  is isomorphic to the direct sum of  $(V_X \otimes \mathcal{O}_X)/\mathcal{O}_X(-1)$  with a trivial bundle.*

*Proof.* Case-by-case verification using the explicit form of the embeddings given in Theorems 2.4 and 2.5.  $\square$

Let us illustrate how the subbundle  $\varphi^*\mathcal{S}_Y \subset V_s \otimes \mathcal{O}_X$  recovers the linear embedding  $X \xrightarrow{\varphi} Y$ . For instance, assume that  $X \xrightarrow{\varphi} Y$  is a combination of standard and isotropic extensions

$$X = GO(m, V_n) \xrightarrow{\tau_X} G(m, V_n) \xrightarrow{\sigma} G(l, V_r) \xrightarrow{\iota} GO(l, V_s) = Y ,$$

$\sigma$  being a non-strict standard extension given by  $\sigma(U) = U^\perp \oplus W$ , where  $U^\perp \subset V_n^*$ ,  $W \subset V_r$  is a fixed subspace together with a monomorphism  $V_n^* \oplus W \hookrightarrow V_r$ . The isotropic extension  $\iota$  turns  $V_n^* \oplus W$  into an isotropic subspace of the orthogonal space  $V_s$ .

We have  $\varphi^*\mathcal{S}_Y \cong \mathcal{S}_X^\perp \oplus (W \otimes \mathcal{O}_X)$ . To read back  $\varphi$  from the monomorphism

$$(4) \quad \varphi^*\mathcal{S}_Y \cong \mathcal{S}_X^\perp \oplus (W \otimes \mathcal{O}_X) \hookrightarrow V_s \otimes \mathcal{O}_X ,$$

observe that simply

$$(5) \quad \varphi(U) = (\varphi^*\mathcal{S}_Y)_U$$

where  $U \in X$  and  $(\varphi^*\mathcal{S}_Y)_U$  denotes the geometric fibre of the bundle  $\varphi^*\mathcal{S}_Y$  the point  $U$ . It is the morphism (4), not just the bundle  $\varphi^*\mathcal{S}_Y$ , which ensures that the map (5) coincides with  $\varphi$ . In particular, note that  $\iota(V_n^* \oplus W)$  is the union of the images in  $V_s$  of all geometric fibres of  $\varphi^*\mathcal{S}_Y$ , and the space  $W$  is the intersection of all such images.

If  $\varphi$  instead equals a mixed combination of standard and isotropic extensions of the form

$$X = GO(m, V_n) \xrightarrow{\tau_X} G(m, V_n) \xrightarrow{\sigma} G(l, V_r) \xrightarrow{\iota} GS(l, V_s) = Y ,$$

then the same holds with the only change that the form on  $V_s$  is now symplectic.

### 3. SPECIAL AND MAXIMAL LINEAR EMBEDDINGS

In this section we complete the classification of linear embeddings of Grassmannians. As a corollary we classify maximal linear embeddings of Grassmannians.

**3.1. Embeddings of Spinor Grassmannians.** In this subsection we classify embeddings of spinor Grassmannians into arbitrary Grassmannians. However, we start by defining certain linear embeddings of Grassmannians into a spinor Grassmannian.

Let  $X := GO(m, V_{2m})$  for  $m \geq 5$ . Fix  $U_m \in X$  and set  $V_m := U_m$ . For every splitting  $V_{2m} = U_m \oplus U_m^*$  of  $V_{2m}$  into a sum of maximal isotropic subspaces, and any  $1 \leq k \leq (m-1)/2$ , we define the embedding

$$(6) \quad \theta_m^{2k} : G(m-2k, V_m) \hookrightarrow GO(m, V_{2m}), \quad U \mapsto U \oplus (U^\perp \cap U_m^*).$$

Next, fix  $\bar{U}_m \in X^- := GO^-(m, V_{2m})$  and set  $V_m := \bar{U}_m$ . For every splitting  $V_{2m} = \bar{U}_m \oplus \bar{U}_m^*$  and any  $1 \leq k \leq (m-1)/2$ , we define the embedding

$$(7) \quad \theta_m^{2k-1} : G(m-2k+1, V_m) \hookrightarrow GO(m, V_{2m}), \quad U \mapsto U \oplus (U^\perp \cap \bar{U}_m^*).$$

The embeddings  $\theta_m^r$ ,  $1 \leq r \leq m-1$ , are linear because they send projective lines on  $G(m-r, V_m)$  to projective lines on  $GO(m, V_{2m})$ , which is verified in a straightforward manner.

With help of the embedding  $\theta_m^2$  we are now able to classify linear embeddings of spinor Grassmannians into non-spinor Grassmannians. Note that the linear embeddings of spinor Grassmannians into spinor Grassmannians are classified in Theorem 2.4,(ii).

**Proposition 3.1.** *Every linear embedding  $GO(m, V_{2m}) \xrightarrow{\varphi} Y$ , for  $Y = G(l, V_s)$  or  $Y = GS(l, V_s)$ , factors through a projective space. Every linear embedding  $GO(m, V_{2m}) \xrightarrow{\varphi} Y = GO(l, V_s)$ , where  $s \neq 2l, 2l+1, 2l+2$ , factors through a projective space or a standard quadric on  $Y$ .*

*Proof.* Assume first that  $Y = G(l, V_s)$ . Fix  $V_m \in GO(m, V_{2m})$  and let  $\theta := \theta_m^2$ . We consider the composition

$$G(m-2, V_m) \xrightarrow{\theta} GO(m, V_{2m}) \xrightarrow{\varphi} G(l, V_s).$$

By Theorem 2.4, this composition is either a standard extension or factors through a projective space on  $G(l, V_s)$ .

We claim that in fact  $\varphi \circ \theta$  factors through a projective space. To see this, fix an arbitrary point  $y \in G(m-2, V_m)$  and its images  $\theta(y) \in GO(m, V_{2m})$  and  $\varphi \circ \theta(y) \in G(l, V_s)$ . Recall from Subsection 1.3 that in each of the Grassmannians  $G(m-2, V_m)$ ,  $GO(m, V_{2m})$ , and  $G(l, V_s)$  there are two families of maximal projective spaces passing through the respective points  $y, \theta(y), \varphi \circ \theta(y)$ . In  $GO(m, V_{2m})$  one family consists of  $\mathbb{P}^{m-1}$ -s, the other of  $\mathbb{P}^3$ -s, and every space of the first family intersects every space of the second family in a projective plane. On the other hand, in both Grassmannians  $G(m-2, V_m)$  and  $G(l, V_s)$  any two maximal projective spaces from different families intersect in a projective line. This implies that  $\varphi$  cannot separate the two families of maximal projective spaces on  $GO(m, V_{2m})$  passing through  $\theta(y)$ , in the sense of sending two spaces from different families on  $GO(m, V_{2m})$  to two respective spaces on  $G(l, V_s)$  also belonging to different families. Since any standard extension  $G(m-2, V_m) \hookrightarrow G(l, V_s)$  separates the two families of maximal projective spaces on  $G(m-2, V_m)$ , we conclude that the composition  $\varphi \circ \theta$  is not a standard extension. Hence its image  $\varphi \circ \theta(G(m-2, V_m))$  lies in some maximal projective space  $\mathbb{P}_0^N$  in  $G(l, V_s)$  containing the point  $\varphi \circ \theta(y)$ .

The space  $\mathbb{P}_0^N$  intersects any other maximal projective space from its own family in  $G(l, V_s)$  only at the point  $\varphi \circ \theta(y)$ . Any maximal  $\mathbb{P}^3$  passing through  $\varphi \circ \theta(y)$  in  $GO(m, V_{2m})$  intersects  $\mathbb{P}_0^N$  along a copy of  $\mathbb{P}^1$ , so all such  $\mathbb{P}^3$ -s must be contained in  $\mathbb{P}_0^N$ . The same argument shows that  $\mathbb{P}_0^N$  contains all maximal  $\mathbb{P}^{m-1}$ -s on  $GO(m, V_{2m})$  passing through  $\theta(y)$ . Hence all projective lines on  $GO(m, V_{2m})$  passing through  $\theta(y)$  are contained in  $\mathbb{P}_0^N$ . By Lemma 1.2 any two points in  $GO(m, V_{2m})$  are connected via a finite chain of projective lines, and the fact that the point  $y$  is arbitrary enables us to complete the argument for  $Y = G(l, V_s)$ .

The cases where  $Y$  is isomorphic to  $GS(l, V_s)$  or  $GO(l, V_s)$  are deduced from the case of  $G(l, V_s)$  by an argument analogous to the one used in the proof of Theorem 2.5,(v). Namely,

we compose  $\varphi$  with the tautological embedding  $\tau_Y : Y \hookrightarrow G(l, V_s)$  and use the fact that the embedding  $\tau_Y \circ \varphi : GO(m, V_{2m}) \hookrightarrow G(l, V_s)$  factors through a maximal projective space  $P$  on  $G(l, V_s)$ . If  $Y = GS(l, V_s)$  then the intersection  $P \cap \tau_Y(Y)$  is a projective space on  $Y$  containing the image of  $GO(m, V_{2m})$ , so the statement holds in this case. If  $Y = GO(l, V_s)$  then  $P \cap \tau_Y(Y)$  is either a projective space or a maximal standard quadric  $Q$  on  $Y$ . Hence, if  $GO(m, V_{2m}) \xrightarrow{\varphi} Y$  does not factor through a projective space then  $\varphi$  factors through a standard quadric on  $Y$ , and the embedding of  $GO(m, V_{2m})$  into this quadric is one of the embeddings from Proposition 2.1. This completes the proof.  $\square$

**3.2. The Grassmannians  $GO(n-2, V_{2n})$ .** We now classify linear embeddings of  $GO(n-2, V_{2n})$  into non-spinor Grassmannians. Set  $X := GO(n-2, V_{2n})$  throughout this subsection. First we construct a special linear embedding of  $X$  into  $G(2, V_{2n-1})$ .

Let  $S := GO(n, V_{2n})$ . The vector space  $V_S := H(S, \mathcal{O}_S(1))^*$  has dimension  $2^{n-1}$ , and

$$\pi_S : S \rightarrow \mathbb{P}(V_S)$$

is a minimal projective embedding of  $S$ . Set  $Y := G(2, V_S) \cong G(2, V_{2n-1})$  and recall that this Grassmannian can be interpreted as the variety of projective lines in  $\mathbb{P}(V_S)$ . Furthermore, from Subsection 1.3.6 we know that  $X$  is the variety of projective lines on  $S$ , and formula (3) allows us to parametrize the projective lines on  $S$  as

$$\mathbb{P}_{S,U}^1 := \{V \in S : U \subset V\} \subset S \quad \text{for } U \in GO(n-2, V_{2n}) = X.$$

**Proposition 3.2.** *The map*

$$(8) \quad \delta_n : GO(n-2, V_{2n}) \rightarrow G(2, V_S), \quad U \mapsto \pi_S(\mathbb{P}_{S,U}^1)$$

*is a linear embedding and its image is the variety of projective lines on  $S$ .*

*Proof.* It is clear that  $\delta_n$  is an injection, and it is routine to check that  $\delta_n$  is an embedding of algebraic varieties. The fact that the image of  $\delta_n$  is exactly the variety of projective lines on  $S$  follows directly from the construction. To prove that  $\delta_n$  is linear we will show that it sends every projective line on  $X$  to a projective line on  $Y$ . A projective line on  $X$  has the form

$$\mathbb{P}_{X, U_{n-3} \subset U_{n-1}}^1 = \{U \in X : U_{n-3} \subset U \subset U_{n-1}\}$$

for some fixed isotropic subspaces  $U_{n-3} \subset U_{n-1} \subset V_{2n}$  of the indicated dimensions. The points on  $\mathbb{P}_{X, U_{n-3} \subset U_{n-1}}^1$  correspond precisely to those projective lines in  $S$  which contain the unique maximal isotropic subspace  $U_n \subset V_{2n}$  satisfying  $U_{n-1} \subset U_n \in S$ .

The union of these projective lines on  $S$  is the following projective plane on  $S$ :

$$\mathbb{P}_{S, U_{n-3} \subset \bar{U}_n}^2 = \{V \in GO(n, V_{2n}) : U_{n-3} \subsetneq (V \cap U_{n-1})\} = \bigcup_{U \in \mathbb{P}_{X, U_{n-3} \subset U_{n-1}}^1} \mathbb{P}_{S,U}^1,$$

where  $\bar{U}_n \subset V_{2n}$  is the unique maximal isotropic subspace such that  $U_n \cap \bar{U}_n = U_{n-1}$ . (Note that  $\bar{U}_n \notin S$ .) It follows that the image of  $\mathbb{P}_{X, U_{n-3} \subset U_{n-1}}^1$  in  $Y$  under  $\delta_n$  is given by the variety of projective lines on the projective plane  $\mathbb{P}_{S, U_{n-3} \subset \bar{U}_n}^2$  containing the point  $U_n$ . Since this variety is a projective line on  $Y$  the proof is complete.  $\square$

Recall that the spinor Grassmannian  $S$  used to define the embedding  $\delta_n$  is a connected component of the variety of maximal isotropic subspaces of  $V_{2n}$ . If one uses the other connected component  $S^-$  of the same variety, one obtains an embedding  $GO(n-2, V_{2n}) \hookrightarrow G(2, V_{S^-})$  which is a composition of an isomorphism  $G(2, V_S) \xrightarrow{\sim} G(2, V_{S^-})$  with the embedding  $\delta_n$ .

Let  $\mathcal{S}_2 := \mathcal{S}_{G(2, V_S)}$  be the tautological bundle on  $G(2, V_S)$ . The bundle  $\delta_n^* \mathcal{S}_2$  is the  $SO(V_{2n})$ -linearized vector bundle of rank 2 whose fibre at  $U \in X$  is the 2-dimensional space  $\pi_S(\mathbb{P}_{S,U}^1)$ .

The embedding  $\delta_n$  factors through the tautological embedding of some orthogonal or symplectic Grassmannian in  $G(2, V_S)$  whenever  $V_S$  admits a respective non-degenerate bilinear form for which all two dimensional subspaces of  $V_S$  corresponding to projective lines on  $S$  are isotropic.

If  $\delta_n$  factors through  $GO(2, V_S)$  (respectively,  $GS(2, V_S)$ ) we denote by  $\delta_n^O$  (respectively,  $\delta_n^S$ ) the embedding such that  $\delta_n = \tau \circ \delta_n^O$  (respectively,  $\delta_n = \tau \circ \delta_n^S$ ):

$$(9) \quad \delta_n : GO(n-2, V_{2n}) \xrightarrow{\delta_n^O} GO(2, V_S) \xrightarrow{\tau} G(2, V_S),$$

$$(10) \quad \delta_n : GO(n-2, V_{2n}) \xrightarrow{\delta_n^S} GS(2, V_S) \xrightarrow{\tau} G(2, V_S).$$

In fact, if  $n$  is even then Proposition 4.4 below implies that  $\delta_n$  always factors through at least one of  $GO(2, V_S)$  and  $GS(2, V_S)$ .

The following theorem is our main result in this subsection.

**Theorem 3.3.** *The following statements hold for every  $n \geq 4$ .*

- (i) *Every linear embedding  $GO(n-2, V_{2n}) \hookrightarrow G(l, V_r)$  factors through a projective space, or factors through the tautological embedding and a standard extension  $\sigma$  as*

$$GO(n-2, V_{2n}) \xrightarrow{\tau_X} G(n-2, V_{2n}) \xrightarrow{\sigma} G(l, V_r),$$

*or factors through the embedding  $\delta_n$  and a standard extension  $\sigma'$  as*

$$GO(n-2, V_{2n}) \xrightarrow{\delta_n} G(2, V_{2n-1}) \xrightarrow{\sigma'} G(l, V_r).$$

- (ii) *Every linear embedding  $GO(n-2, V_{2n}) \xrightarrow{\varphi} GO(l, V_r)$ ,  $l < \lfloor \frac{r-1}{2} \rfloor$ , is a standard extension, or factors through a standard quadric, or factors as*

$$GO(n-2, V_{2n}) \xrightarrow{\psi} G(l, V_{\lfloor r/2 \rfloor}) \xrightarrow{\iota} GO(l, V_r)$$

*where  $\iota$  is an isotropic extension and  $\psi$  is one of the embeddings of part (i), or factors as*

$$GO(n-2, V_{2n}) \xrightarrow{\delta_n^O} GO(2, V_{2n-1}) \xrightarrow{\sigma} GO(l, V_r)$$

*for a standard extension  $\sigma$ .*

- (iii) *Every linear embedding  $GO(n-2, V_{2n}) \xrightarrow{\varphi} GS(l, V_{2r})$  factors as*

$$GO(n-2, V_{2n}) \xrightarrow{\psi} G(l, V_r) \xrightarrow{\iota} GS(l, V_{2r})$$

*where  $\iota$  is an isotropic extension and  $\psi$  is one of the embeddings of part (i), or factors as*

$$GO(n-2, V_{2n}) \xrightarrow{\delta_n^S} GS(2, V_{2n-1}) \xrightarrow{\sigma} GS(l, V_{2r}).$$

*for a standard extension  $\sigma$ .*

*Proof.* We assume first  $n \geq 5$ . For part (i), let  $\varphi : X \hookrightarrow Y = G(l, V_r)$  be a linear embedding which does not factor through a projective space. We borrow the strategy of [9] and consider the images of maximal standard quadrics on  $X$  under  $\varphi$ . Such a maximal quadric is 4-dimensional, has the form  $Q := Q_{X, U_{n-3}}^4$  for some  $U_{n-3} \in GO(n-3, V_{2n})$ , and is isomorphic to the ordinary Grassmannian  $G(2, V_4)$ . There are two options for the restriction  $\varphi|_Q$ :

- 1)  $\varphi|_Q$  factors through a projective space.
- 2)  $\varphi|_Q$  is a standard extension.

In fact, exactly one of these two options holds simultaneously for all maximal standard quadrics on  $X$ . This is due to the irreducibility of the family of maximal standard quadrics in  $X$  and the existence of two irreducible families of maximal projective spaces on  $Y$ ; a detailed argument is given in the proof of [9, Proposition 3.15].

In case 1) the situation is analogous to the one considered in Theorem 2.4, (iii), and we deduce that  $\varphi$  factors through the tautological embedding  $\tau_X$  as  $\varphi = \sigma \circ \tau_X$ , where  $\sigma$  is a standard extension of ordinary Grassmannians.

We suppose now that 2) holds, so that  $\varphi|_Q$  is a standard extension for every maximal standard quadric  $Q$  on  $X$ . We will prove that  $\varphi$  factors through  $\delta_n : X \hookrightarrow G(2, V_{2n-1})$ . For this we need an auxiliary construction.

Any projective line  $L := \mathbb{P}_{X, U_{n-3} \subset U_{n-1}}^1$  on  $X$  is equal to the intersection of three uniquely determined maximal projective spaces on  $X$ , namely,

$$\mathbb{P}_{X, U_{n-3} \subset U_{n-1}}^1 = \mathbb{P}_{X, U_{n-1}}^{n-2} \cap \mathbb{P}_{X, U_{n-3} \subset U_n}^2 \cap \mathbb{P}_{X, U_{n-3} \subset \bar{U}_n}^2$$

where  $U_n \in S$  and  $\bar{U}_n \in S^-$  form the unique pair of maximal isotropic subspaces of  $V_{2n}$  containing  $U_{n-1}$ . Moreover,  $L$  equals the intersection of any two of the above three maximal projective spaces on  $X$ . The spaces  $\mathbb{P}_{X, U_{n-3} \subset U_n}^2$  and  $\mathbb{P}_{X, U_{n-3} \subset \bar{U}_n}^2$  are maximal projective spaces on the standard quadric  $Q_{X, U_{n-3}}^4 \subset X$ . For any  $U \in S \cup S^-$  we denote by  $Z_U \subset X$  the image of the isotropic extension  $G(n-2, U) \hookrightarrow GO(n-2, V_{2n}) = X$ . Then  $\mathbb{P}_{X, U_{n-1}}^{n-2}$  and  $\mathbb{P}_{X, U_{n-3} \subset U_n}^2$  are maximal projective spaces on  $Z_{U_n}$ , while  $\mathbb{P}_{X, U_{n-1}}^{n-2}$  and  $\mathbb{P}_{X, U_{n-3} \subset U_n}^2$  are such on  $Z_{\bar{U}_n}$ . We have  $\mathbb{P}_{X, U_{n-1}}^{n-2} = Z_{U_n} \cap Z_{\bar{U}_n}$ .

The image of  $L$  under  $\varphi$  has the form

$$\varphi(L) = \mathbb{P}_{Y, W_{l-1} \subset W_{l+1}}^1 = \mathbb{P}_{Y, W_{l-1}}^{r-l} \cap \mathbb{P}_{Y, W_{l+1}}^l$$

for a unique flag of subspaces  $W_{l-1} \subset W_{l+1} \subset V_r$ . The image under  $\varphi$  of each of three spaces  $\mathbb{P}_{X, U_{n-1}}^{n-2}$ ,  $\mathbb{P}_{X, U_{n-3} \subset U_n}^2$  and  $\mathbb{P}_{X, U_{n-3} \subset \bar{U}_n}^2$  is contained in exactly one of the subspaces  $\mathbb{P}_{Y, W_{l-1}}^{r-l}$  and  $\mathbb{P}_{Y, W_{l+1}}^l$ . Thus at least two of the three spaces are mapped to the same maximal projective space on  $Y$ . Consequently, the image under  $\varphi$  of at least one of the three Grassmannians  $Q_{X, U_{n-3}}^4$ ,  $Z_{U_n}$ ,  $Z_{\bar{U}_n}$  is contained in a projective space on  $Y$ . If all three Grassmannians are mapped to the same maximal projective space on  $Y$  then  $\varphi$  factors through a projective space on  $Y$ . This case is excluded by our hypothesis, so exactly one of  $Q_{X, U_{n-3}}^4$ ,  $Z_{U_n}$ ,  $Z_{\bar{U}_n}$  is mapped by  $\varphi$  into a projective space on  $Y$ . The possibility that  $\varphi$  maps  $Q_{X, U_{n-3}}^4$  to a projective space is also excluded, because it was considered as case 1). Hence the restriction of  $\varphi$  to one of the Grassmannians  $Z_{U_n}$  and  $Z_{\bar{U}_n}$  is a standard extension, while the restriction to the other Grassmannian factors through a projective space. Up to change of notation, we may assume that  $\varphi|_{Z_{\bar{U}_n}}$  is a standard extension.

To summarize, it remains to consider the case where  $\varphi|_{Q_{X, U_{n-3}}^4}$  and  $\varphi|_{Z_{\bar{U}_n}}$  are standard extensions while  $\varphi|_{Z_{U_n}}$  factors through a projective space. By the irreducibility of the parameter spaces, this property holds for every  $U_{n-3} \in GO(n-3, V_{2n})$ , every  $U_n \in S$  and every  $\bar{U}_n \in S^-$ . In this setting we will construct a linear embedding  $\pi$  of the spinor Grassmannian  $S$  into the projective space  $\mathbb{P}(V_r^*)$ , which will help us see that  $\varphi$  factors through  $\delta_n$ .

For every  $U_n \in S$  the image  $\varphi(Z_{U_n})$  is contained in a maximal projective space on  $Y$  of the form  $\mathbb{P}_{Y, W_{l+1}}^l$  or  $\mathbb{P}_{Y, W_{l-1}}^{r-l}$ . These two cases are analogous and interchangeable by the use of the isomorphism  $G(l, V_r) \cong G(r-l, V_r^*)$ . We assume  $\varphi(Z_{U_n}) \subset \mathbb{P}_{Y, W_{l+1}}^l$ , which yields a morphism

$$\pi : S \rightarrow G(l+1, V_r), \quad U_n \mapsto W_{l+1}.$$

On the other hand, for every  $\bar{U}_n \in S^-$  the restriction of  $\varphi$  to  $Z_{\bar{U}_n} = G(n-2, \bar{U}_n)$  is a standard extension which is strict because of the above choice of  $\mathbb{P}_{Y, W_{l+1}}^l$  instead of  $\mathbb{P}_{Y, W_{l-1}}^{r-l}$ . Thus there is an inclusion  $\nu_{\bar{U}_n} : \bar{U}_n \hookrightarrow V_r$  and a subspace  $W_{\bar{U}_n} \subset V_r$  of dimension  $l+2-n$  such that

$$\varphi(U) = W_{\bar{U}_n} \oplus \nu_{\bar{U}_n}(U)$$

for  $U \in Z_{\bar{U}_n}$ . We obtain a morphism

$$\rho : S^- \rightarrow G(l+2, V_r), \quad \bar{U}_n \mapsto W_{\bar{U}_n} \oplus \nu_{\bar{U}_n}(\bar{U}_n).$$

For  $U_n \in S$  and  $\bar{U}_n \in S^-$  the intersection  $Z_{U_n} \cap Z_{\bar{U}_n}$  in  $X$  is empty or is a maximal projective space on  $X$  of the form  $\mathbb{P}_{X, U_{n-1}}^{n-2}$  whenever  $U_{n-1} = U_n \cap \bar{U}_n$  has dimension  $n-1$ . Therefore,

there is a third morphism

$$\pi' : G(n-1, V_{2n}) \rightarrow G(l+1, V_r), \quad U_{n-1} \rightarrow \pi(U_n) = W_{\bar{U}_n} \oplus \nu_{\bar{U}_n}(U_{n-1}),$$

where  $U_n \in S$  and  $\bar{U}_n \in S^-$  is the unique pair such that  $U_n \cap \bar{U}_n = U_{n-1}$ . We note that  $\pi$  and  $\pi'$  have the same image, and  $\pi(U_n) = \pi'(U_{n-1})$  if and only if  $U_{n-1} \subset U_n$ .

We claim that  $\rho$  is a constant morphism and  $\pi$  is an linear embedding.

To prove that the morphism  $\rho$  is constant it suffices to show that it is constant on any projective line on  $S^-$ . Fix  $\bar{U}_n, \bar{U}'_n \in S$  such that  $\bar{U}_n \cap \bar{U}'_n =: U_{n-2} \in X$ , so that  $\bar{U}_n, \bar{U}'_n$  are two distinct points on the projective line  $\mathbb{P}_{S^-, U_{n-2}}^1$ . The element  $U_{n-2}$  also determines a projective line on  $S$ , and we fix  $U_n, U'_n \in \mathbb{P}_{S, U_{n-2}}^1$  such that  $U_n \cap U'_n = U_{n-2}$ . Then each of the intersections

$$U_{n-1}^{(1)} := U_n \cap \bar{U}_n, \quad U_{n-1}^{(2)} := U'_n \cap \bar{U}_n, \quad U_{n-1}^{(3)} := U_n \cap \bar{U}'_n, \quad U_{n-1}^{(4)} := U'_n \cap \bar{U}'_n$$

has dimension  $n-1$  and contains  $U_{n-2}$ . Hence we have

$$\varphi(U_{n-2}) \subset \pi(U_n) \cap \pi(U'_n), \quad \pi(U_n) + \pi(U'_n) \subset \rho(\bar{U}_n) \cap \rho(\bar{U}'_n).$$

Each of  $\pi(U_n)$  and  $\pi(U'_n)$  has codimension 1 in each of  $\rho(\bar{U}_n)$  and  $\rho(\bar{U}'_n)$ , therefore

$$\rho(\bar{U}_n) = \rho(\bar{U}'_n) \iff \pi(U_n) \neq \pi(U'_n) \iff \pi(U_n) \cap \pi(U'_n) = \varphi(U_{n-2}).$$

In other words, the restriction of  $\rho$  to  $\mathbb{P}_{S^-, U_{n-2}}^1$  is constant if and only if the restriction of  $\pi$  to  $\mathbb{P}_{S, U_{n-2}}^1$  is a linear embedding.

Since  $U_{n-1}^{(1)}$  and  $U_{n-1}^{(2)}$  are distinct hyperplanes in  $\bar{U}_n$  they define distinct maximal projective spaces on  $Z_{\bar{U}_n}$ . It follows that  $\pi'(U_{n-1}^{(1)}) \neq \pi'(U_{n-1}^{(2)})$  because any standard extension of ordinary Grassmannians maps distinct maximal projective spaces on its domain to distinct maximal projective spaces on the target Grassmannian. Hence

$$\pi(U_n) = \pi'(U_{n-1}^{(1)}) \neq \pi'(U_{n-1}^{(2)}) = \pi(U'_n).$$

We can conclude that the restriction of  $\pi$  to every projective line on  $S$  is a linear embedding, and  $\rho$  is a constant morphism.

Since the restriction of  $\pi$  to any projective line is linear, the map  $\pi$  is itself linear. To prove that  $\pi$  is a linear embedding it remains to show that it is injective. Let  $V_{l+2} := \rho(S^-) \in G(l+2, V_r)$  be the image of  $\rho$ . For every  $\bar{U}_n \in S^-$  we have  $\varphi(Z_{\bar{U}_n}) \subset \sigma_1(G(l, V_{l+2})) \subset Y$ , where  $\sigma_1 : G(l, V_{l+2}) \hookrightarrow G(l, V_r)$  is the canonical standard extension. It follows that the embedding  $\varphi$  factors through the standard extension  $\sigma_1$ , i.e., there exists a linear embedding  $\psi : X \rightarrow Z := G(l, V_{l+2})$  such that  $\varphi = \sigma_1 \circ \psi$ . This allows us to reduce to the case where  $V_r = V_{l+2}$  and  $\varphi = \psi$ .

Suppose  $\pi$  is not injective. In other words,  $\pi(U_n) = \pi(U''_n) =: W_{l+1}$  for some  $U_n, U''_n \in S$  which do not belong to a projective space on  $S$ . Then  $\dim(U_n \cap U''_n) = n-2k$  for some  $k > 1$ . Let  $U_{n-2}, U''_{n-2} \in X$  be such that  $U_{n-2} \subset U_n$ ,  $U''_{n-2} \subset U''_n$  and  $U_n \cap U''_n = U_{n-2} \cap U''_{n-2}$ . The images  $\varphi(U_{n-2})$  and  $\varphi(U''_{n-2})$  are then two distinct hyperplanes in  $\pi(U_n)$ . Hence the intersection  $W_{l-1} := \varphi(U_{n-2}) \cap \varphi(U''_{n-2})$  has dimension  $l-1$ , and we deduce that the two images  $\varphi(U_{n-2})$  and  $\varphi(U''_{n-2})$  belong to the projective line  $\mathbb{P}_{Y, W_{l-1} \subset W_{l+1}}^1$  on  $Y = G(l, V_{l+2})$ . Since  $\varphi$  is a linear embedding,  $U_{n-2}$  and  $U''_{n-2}$  belong to a projective line  $\mathbb{P}_{X, U_{n-3} \subset U_{n-1}}^1$  on  $X$ . In particular  $\dim(U_{n-2} \cap U''_{n-2}) = n-3$ , a contradiction.

This enables us to conclude that  $\pi$  is a linear embedding, and we can define the previously announced embedding  $\underline{\pi}$  as the composition of  $\pi$  with the duality isomorphism

$$\underline{\pi} : S \xrightarrow{\pi} G(l+1, V_{l+2}) \cong \mathbb{P}(V_{l+2}^*).$$

Similarly, let  $\underline{\varphi}$  be the composition

$$\underline{\varphi} : X \xrightarrow{\varphi} G(l, V_{l+2}) \cong G(2, V_{l+2}^*).$$

The explicit form of the embedding  $\underline{\pi}$  yields immediately to our desired factorization property of the embedding  $\varphi$ . Indeed,  $\underline{\pi} : S \rightarrow V_{l+2}^*$  induces a linear map  $\hat{\underline{\pi}} : V_S \hookrightarrow V_{l+2}^*$ , which in turn induces a standard extension  $\sigma_2 : G(2, V_S) \hookrightarrow G(2, V_{l+2}^*)$  through which  $\underline{\varphi}$  factors. The resulting embedding  $X \hookrightarrow G(2, V_S)$  is

$$\underline{\varphi}(U_{n-2}) = \underline{\pi}(\mathbb{P}_{S, U_{n-2}}^1) = \mathbb{P}(\underline{\pi}(U_n) + \underline{\pi}(U)) = \mathbb{P}(\underline{\varphi}(U_{n-2})) \in \sigma_2(G(2, V_S)) ,$$

which is identical with  $\delta_n$  by (8). Therefore  $\underline{\varphi} = \sigma_2 \circ \delta_n$ , and the initial embedding  $\varphi$  equals the composition

$$\varphi : X \xrightarrow{\delta_n} G(2, V_S) \xrightarrow{\sigma_2} G(2, V_{l+2}^*) \cong G(l, V_{l+2}) \xrightarrow{\sigma_1} G(l, V_r) = Y .$$

This completes the proof of part (i) for  $n \geq 5$ .

It remains to consider the case  $n = 4$ . Here the embedding  $\delta_4$  is related to the tautological embedding  $\tau_X$  via the isomorphism  $GO(1, V_8) \cong GO(4, V_8)$ , and the two factorizations in part (i) are the same. The result is that every linear embedding  $GO(2, V_8) \hookrightarrow Y$ , where  $Y$  is an ordinary or a symplectic Grassmannian, factors through the tautological embedding into  $G(2, V_8)$ . In case  $Y \cong GO(l, V_r)$ , there are also the options for the embedding to be a standard extension and to factor through a standard quadric. This settles parts (i),(ii),(iii) for  $n = 4$ .

Parts (ii) and (iii) for  $n \geq 5$  follow from part (i) by use of the composition  $\tau_Y \circ \varphi$  of a given embedding  $\varphi$  of  $X$  into an isotropic Grassmannian  $Y$  with the tautological embedding  $\tau_Y$  of  $Y$ .  $\square$

**3.3. The Grassmannians  $GO(n-1, V_{2n+1})$ .** In this subsection we classify linear embeddings of the Grassmannian  $X := GO(n-1, V_{2n+1})$  for  $n \geq 3$  into non-spinor Grassmannians. This case is similar and closely related to the case of  $GO(n-2, V_{2n})$  considered in the previous subsection.

There is a special linear embedding of  $X$ , obtained as the composition

$$(11) \quad \delta_{n+1}^1 : GO(n-1, V_{2n+1}) \xrightarrow{\sigma_1} GO(n-1, V_{2n+2}) \xrightarrow{\delta_{n+1}^1} G(2, V_{2n}) = G(2, V_{GO(n, V_{2n+1})})$$

of a standard extension  $\sigma_1$  and the embedding  $\delta_{n+1}$  given in (8). If the embedding  $\delta_{n+1}^1$  factors through a tautological embedding  $GO(2, V_{2n}) \hookrightarrow G(2, V_{2n})$  or  $GS(2, V_{2n}) \hookrightarrow G(2, V_{2n})$ , we can write

$$(12) \quad \begin{aligned} \delta_{n+1}^1 : G(n-1, V_{2n+1}) &\xrightarrow{\delta_{n+1}^{1,O}} GO(2, V_{2n}) \hookrightarrow G(2, V_{2n}) , \\ \delta_{n+1}^1 : G(n-1, V_{2n+1}) &\xrightarrow{\delta_{n+1}^{1,S}} GS(2, V_{2n}) \hookrightarrow G(2, V_{2n}) . \end{aligned}$$

for respective embeddings  $\delta_{n+1}^{1,O}$  and  $\delta_{n+1}^{1,S}$ . Proposition 4.4 below implies that one of these factorizations always exists.

**Theorem 3.4.** *The following statements hold for  $n \geq 3$ .*

(i) *Every linear embedding  $GO(n-1, V_{2n+1}) \hookrightarrow G(l, V_r)$  factors as*

$$(13) \quad GO(n-1, V_{2n+1}) \xrightarrow{\tau_X} G(n-1, V_{2n+1}) \xrightarrow{\sigma} G(l, V_r)$$

*or as*

$$GO(n-1, V_{2n+1}) \xrightarrow{\delta_{n+1}^1} G(2, V_{2n}) \xrightarrow{\sigma'} G(l, V_r) ,$$

*where  $\sigma$  and  $\sigma'$  are standard extensions, or factors through a projective space.*

(ii) *Every linear embedding  $GO(n-1, V_{2n+1}) \hookrightarrow GO(l, V_r)$ ,  $l \leq \lfloor \frac{r-1}{2} \rfloor$ , is a standard extension, or factors through a standard quadric, or factors as*

$$GO(n-1, V_{2n+1}) \xrightarrow{\psi} G(l, V_k) \xrightarrow{\iota} GO(l, V_r)$$

where  $\psi$  is one of the embeddings in part (i) and  $\iota$  is an isotropic extension, or factors as

$$GO(n-1, V_{2n+1}) \xrightarrow{\delta_{n+1}^{1,O}} GO(2, V_{2n}) \xrightarrow{\sigma'} GO(l, V_r)$$

where  $\sigma, \sigma'$  are standard extensions and  $\delta_{n+1}^{1,O}$  is the embedding given in (12).

(iii) Every linear embedding  $GO(n-1, V_{2n+1}) \hookrightarrow GS(l, V_r)$  factors as

$$GO(n-1, V_{2n+1}) \xrightarrow{\psi} G(l, V_k) \xrightarrow{\iota} GS(l, V_r)$$

where  $\psi$  is one of the embeddings in part (i) and  $\iota$  is an isotropic extension, or factors as

$$GO(n-1, V_{2n+1}) \xrightarrow{\delta_{n+1}^{1,S}} GS(2, V_{2n}) \xrightarrow{\sigma'} GS(l, V_r)$$

where  $\sigma, \sigma'$  are standard extensions and  $\delta_{n+1}^{1,S}$  is the embedding given in (12).

*Proof.* The proof follows the same lines as the proof of Theorem 3.3. We outline the main steps for part (i). Let  $\varphi : X := GO(n-1, V_{2n+1}) \hookrightarrow G(l, V_r) =: Y$  be a linear embedding. A maximal standard quadric  $Q$  on  $X$  is 3-dimensional and has the form  $Q = Q_{X, U_{n-2}}^3$  given in Subsection 1.3.5. Any 3-dimensional quadric is isomorphic to  $GS(2, V_4)$ . Hence there are the following two options for the restriction of  $\varphi$  to  $Q$ :

1)  $\varphi|_Q$  factors through a projective space.

2)  $\varphi|_Q$  factors through a composition  $\sigma \circ \tau$  where  $\tau$  is the tautological embedding of  $GS(2, V_4)$  and  $\sigma : G(2, V_4) \rightarrow Y$  is a standard extension.

If 1) holds one can show that the embedding  $\varphi$  factors through the tautological embedding  $\tau_X : GO(n-1, V_{2n+1}) \hookrightarrow G(n-1, V_{2n+1})$ . Thus  $\varphi = \sigma \circ \tau_X$  where  $\sigma : G(n-1, V_{2n+1}) \hookrightarrow G(l, V_r)$  is a linear embedding. By Theorem 2.4,  $\sigma$  is a standard extension or factors through a projective space, and we obtain a factorization of the type (13).

If 2) holds we claim that the embedding  $\varphi$  factors through the embedding  $\delta_{n+1}^1$ . To show this we construct an auxiliary linear embedding  $\pi : S \rightarrow \mathbb{P}(V_r)$  of the spinor Grassmannian  $S := GO(n, V_{2n+1})$ . The embedding  $\pi$  is constructed first for  $l = 2$ , using the fact that  $S$  is the parameter space for the family of maximal projective spaces on  $X$  (see Subsection 1.3.5). Then one shows that  $\varphi$  factors through a standard extension  $\sigma_1 : X \hookrightarrow GO(n-1, V_{2n+2})$  using the isomorphism of spinor Grassmannians  $S \cong GO(n+1, V_{2n+2}) =: S'$  and the identification of  $GO(n-1, V_{2n+2})$  with the variety of projective lines on  $S$  given by the embedding  $\delta_{n+1}^1$ . Here  $V_{2n+2} := V_{2n+1} \oplus V_1$  is endowed with a non-degenerate symmetric bilinear form extending the given form on  $V_{2n+1}$ . Thus  $\varphi$  factors through  $\delta_{n+1}^1$ , and hence (i) holds.

To prove parts (ii) and (iii) one can compose any given linear embedding  $X \xrightarrow{\varphi} Y$  into a symplectic or orthogonal non-spinor Grassmannian  $Y$  with the tautological embedding  $\tau_Y$ . Then part (i) applies to  $\tau_Y \circ \varphi$ , and the result follows in a straightforward manner.  $\square$

**3.4. Embeddings into spinor Grassmannians.** We are now ready to classify the linear embeddings of Grassmannians  $X$  into spinor Grassmannians, with the following exceptions:  $X = GS(m, V_{2m})$  for  $m \geq 2$ ,  $X = GO(m-2, V_{2m-\varepsilon})$  for  $5 \leq m$ ,  $\varepsilon \in \{0, 1\}$ .

Let  $X = G(m, V_n), GO(m, V_n), GS(m, V_n)$ , and set respectively

$$X_{\pm 1} := \begin{cases} G(m \pm 1, V_n) & \text{for } m \neq 1, n-1, \\ GO(m \pm 1, V_n) & \text{for } 1 < m < \lfloor n/2 \rfloor, \\ GS(m \pm 1, V_n) & \text{for } 1 < m < n/2. \end{cases}$$

We say that  $X$  is **regular** if both  $X_{-1}$  and  $X_{+1}$  are defined and are non-spinor Grassmannians. An ordinary or symplectic Grassmannian  $X$  is regular if and only if  $X$  has two families of maximal projective spaces parametrized respectively by  $X_{-1}$  and  $X_{+1}$ . If  $X$  is a regular orthogonal Grassmannian then  $X_{+1}$  parametrizes the maximal projective spaces on  $X$  which are not contained in a quadric, while  $X_{-1}$  parametrizes the maximal standard quadrics on  $X$ .

The non-regular Grassmannians are projective spaces, quadrics of dimension at least 5, spinor Grassmannians, and the above mentioned  $GS(m, V_{2m})$  and  $GO(m-2, V_{2m-\varepsilon})$ .

By  $S = S^+ := GO(l, V_{2l})$  we denote a fixed spinor Grassmannian for  $l \geq 4$ . Recall that there are two families of maximal projective spaces on  $S$ : a family of maximal  $\mathbb{P}^{l-1}$ -s parametrized by  $S^- := GO^-(l, V_{2l})$ , and a family of maximal 3-dimensional projective spaces  $\mathbb{P}_{S, W_{l-3}}^3$  parametrized by  $W_{l-3} \in S_{-3} := GO(l-3, V_{2l})$ . Note also that  $\mathbb{P}^3$  is isomorphic to the spinor Grassmannian  $GO(3, V_6)$ , and the inclusion of a maximal  $\mathbb{P}^3$  in  $S$  can be interpreted as a standard extension of spinor Grassmannians.

We first establish several preliminary results needed for the classification given in Theorem 3.10 below.

**Lemma 3.5.** *Let  $Q = Q^{n-2} = GO(1, V_n)$  be an  $n-2$ -dimensional quadric with  $n \geq 7$  and let  $Q \xrightarrow{\varphi} S$  be a linear embedding. Then  $\varphi$  factors through a projective space of the form  $\mathbb{P}_{S, \bar{W}}^{l-1}$  for some  $\bar{W} \in S^-$ , unless  $n \leq 8$  and  $\varphi$  factors through  $Q^6$  as*

$$\varphi : Q \xrightarrow{\sigma_1} Q^6 \cong GO(4, V_8) \xrightarrow{\sigma_2} GO(l, V_{2l}) = S$$

where  $\sigma_1, \sigma_2$  are standard extensions or isomorphisms.

*Proof.* For  $n \geq 10$  the maximal projective spaces on  $Q$  have dimension  $\lfloor n/2 \rfloor - 1 \geq 4$ . Moreover, for any maximal projective space  $P'$  on  $Q$  there is another maximal projective space  $P''$  on  $Q$  such that  $P' \cap P''$  has codimension 1 in both  $P'$  and  $P''$ . On the other hand, no maximal projective spaces on  $S$  intersect in dimension 3 or more. Therefore,  $\varphi$  maps all maximal projective spaces on  $Q$  onto a single projective space  $P$  on  $S$ . For dimension reasons  $P$  is of the form  $\mathbb{P}_{S, \bar{W}}^{l-1}$  for some  $\bar{W} \in S^-$ . Hence  $\varphi$  factors through a projective space. For  $n = 9$  the maximal projective spaces on  $Q$  have dimension 3, and any projective plane in  $Q$  is contained in a  $\mathbb{P}^1$ -family of maximal  $\mathbb{P}^3$ -s. Two maximal projective spaces on  $S$  belonging to a connected family intersect at most at a projective line. Hence  $\varphi$  factors through  $\mathbb{P}_{S, \bar{W}}^{l-1}$  for some  $\bar{W} \in S^-$ .

For  $n = 8$  the quadric  $Q$  is isomorphic to the spinor Grassmannian  $GO(4, V_8)$ . By Theorem 2.4 the embedding  $\varphi$  factors through a projective space or is a standard extension of spinor Grassmannians. For  $n = 7$  the maximal projective spaces of  $Q$  have dimension 2 and are parametrized by the spinor Grassmannian  $GO(3, V_7) \cong GO(4, V_8)$ . Hence for  $U_3 \in GO(3, V_7)$  we have  $\varphi(\mathbb{P}_{Q, U_3}^2) = \mathbb{P}_{S, W_{l-3} \subset \bar{W}_l}^2$  for a unique isotropic flag  $W_{l-3} \subset \bar{W}_l$  in  $V_{2l}$  with  $\bar{W}_l \in S^-$ . This yields morphisms  $\varphi' : GO(3, V_7) \rightarrow S_{-3}$  and  $\varphi'' : GO(3, V_7) \rightarrow S^-$ . If the morphism  $\varphi''$  is constant with value  $\bar{W}_l$  then  $\varphi$  factors through the projective space  $\mathbb{P}_{S, \bar{W}_l}^{l-1}$ . Suppose that  $\varphi''$  is non-constant. Then the image of the linear embedding  $\mathbb{P}(V_7) \hookrightarrow \mathbb{P}(V_S)$  induced by  $\varphi$  intersects  $S$  in  $\varphi(Q)$ . Consequently  $\varphi(Q) \cap \mathbb{P}_{S, W_{l-3}}^3 = \mathbb{P}_{S, W_3 \subset \bar{W}_l}^2$ , so the morphism  $\varphi'$  is injective.

We denote

$$\underline{W} := \bigcap_{U_1 \in Q} \varphi(U_1) = \bigcap_{U_3 \in GO(3, V_8)} \varphi'(U_3)$$

and claim that  $\underline{W}$  has dimension  $l-4$ . Indeed, if  $U_3, U'_3 \in GO(3, V_7)$  is a pair of points such that the corresponding projective planes on  $Q$  intersect along a projective line, then  $U_3 \cap U'_3 =: U_2$  has dimension 2 and hence defines a projective line on  $GO(3, V_7)$  as well as on  $Q$ . We have

$$\varphi(\mathbb{P}_{Q, U_2}^1) = \mathbb{P}_{S, \varphi'(U_3)}^3 \cap \mathbb{P}_{S, \varphi'(U'_3)}^3.$$

Therefore,  $W_{l-2} := \varphi'(U_3) + \varphi'(U'_3)$  is an isotropic subspace of  $V_{2l}$  of dimension  $l-2$ , and  $W_{l-4} := \varphi'(U_3) \cap \varphi'(U'_3)$  has dimension  $l-4$ . Clearly  $\underline{W} \subset W_{l-4}$ , and in fact  $\underline{W} = W_{l-4}$  because any two points on  $GO(3, V_7)$  belong to a connected finite union of projective lines.

The variety  $Z := \{W \in S : \underline{W} \subset W\}$  is the image of a standard extension  $\sigma_2 : GO(4, V_8) \hookrightarrow S$ . Furthermore,  $\varphi$  factors through  $\sigma_2$  and the corresponding embedding  $\sigma_1 : \varphi(Q) \subset Z \cong Q^6$  is necessarily a standard extension of quadrics. The statement follows.  $\square$

**Lemma 3.6.** *Suppose that  $P^{(1)}, P^{(2)}, P^{(3)}, P^{(4)}$  are four projective planes on  $S$  passing through a point  $W_l \in S$ , such that*

$$P^{(1)} \cap P^{(2)} = P^{(3)} \cap P^{(4)} = \{W_l\},$$

*the intersections  $P^{(i)} \cap P^{(j)}$  are projective lines for  $i = 1, 2$  and  $j = 3, 4$ , and none of the unions  $P^{(1)} \cup P^{(2)} \cup P^{(3)}$ ,  $P^{(1)} \cup P^{(2)} \cup P^{(4)}$  is contained in a projective space on  $S$  or in a 6-dimensional quadric on  $S$ . Then  $P^{(3)} \cup P^{(4)}$  is contained in  $\mathbb{P}_{S, \bar{W}_l}^{l-1}$  for some  $\bar{W}_l \in S^-$ .*

*Proof.* Let  $W_{l-3}^{(k)} \subset \bar{W}_l^{(k)}$  be an isotropic flag in  $V_{2l}$  such that  $P^{(k)} = \mathbb{P}_{S, W_{l-3}^{(k)} \subset \bar{W}_l^{(k)}}^2$  for  $k = 1, 2, 3, 4$ .

We aim to show that  $\bar{W}_l^{(3)} = \bar{W}_l^{(4)}$ . Our assumptions imply that the subspaces of  $W_l$  defined below have the indicated dimensions and satisfy the given relations with  $i = 1, 2$  and  $j = 3, 4$ :

$$\begin{aligned} W_{l-4}^{(ij)} &:= W_{l-3}^{(i)} \cap W_{l-3}^{(j)} \subset W_{l-3}^{(i)} + W_{l-3}^{(j)} =: W_{l-2}^{(ij)} \subset W_{l-1}^{(j)} := W_l \cap \bar{W}_l^{(j)}, \\ W_{l-5} &:= W_{l-3}^{(1)} \cap W_{l-3}^{(2)} = W_{l-4}^{(1)} \cap W_{l-4}^{(2)} \subset W_{l-4}^{(1j)} + W_{l-4}^{(2j)} = W_{l-3}^{(j)}, \\ W_{l-3}^{(j)} &= W_{l-2}^{(1j)} \cap W_{l-2}^{(2j)} \subset W_{l-2}^{(1j)} + W_{l-2}^{(2j)} = W_{l-3}^{(1)} + W_{l-3}^{(2)} = W_{l-1}^{(j)}. \end{aligned}$$

In particular  $W_{l-1}^{(3)} = W_{l-1}^{(4)}$ , and hence  $\bar{W}_l^{(3)} = \bar{W}_l^{(4)}$ .  $\square$

**Proposition 3.7.** *Let  $X \xrightarrow{\varphi} S$  be a linear embedding of a regular Grassmannian  $X$  into  $S$ , and assume that  $\varphi$  does not factor through a projective space. Then the image of every maximal projective space on  $X$  is contained in a unique maximal projective space on  $S^-$  of the form  $\mathbb{P}_{S^-, \bar{W}}^{l-1}$  for some  $\bar{W} \in S^-$ . Furthermore,  $\varphi$  induces two linear embeddings*

$$(14) \quad \varphi_{\pm 1} : X_{\pm 1} \rightarrow S^-.$$

*Proof.* Since  $X$  is regular, every maximal projective space on  $X$  has dimension at least 2. Assume that  $X_{+1}$  parametrizes maximal projective spaces  $P$  on  $X$  of dimension  $\dim P \neq 3$ . Then the description of intersections of maximal projective spaces on  $S$  in Subsection 1.3.6 implies that the image  $\varphi(P)$  of any such maximal projective space  $P$  is contained in  $\mathbb{P}_{S, \bar{W}}^{l-1}$  for a unique  $\bar{W} \in S^-$ . This yields a morphism  $\varphi_{+1} : X_{+1} \rightarrow S^-$ ,  $P \mapsto \mathbb{P}_{S, \bar{W}}^{l-1}$ . We will construct the morphism  $\varphi_{-1}$  later, after completing the definition of  $\varphi_{+1}$  in the remaining case.

Suppose that  $X_{+1}$  parametrizes maximal projective spaces of dimension 3 on  $X$ , so  $m = 3$ . Then  $\varphi$  induces a morphism  $\varphi_{+1}$  from  $X_{+1}$  to the variety of all 3-dimensional projective spaces on  $S$ . This variety is disconnected and  $S_{-3}$  is one of its connected components. We have to show that  $\varphi_{+1}(X_{+1}) \subset S^-$ . Assume to the contrary that  $\varphi(\mathbb{P}_{X, U_4}^3) = \mathbb{P}_{S, W_{l-3}}^3$  for some  $U_4 \in X_{+1}$  and  $W_{l-3} \in S_{-3}$ . This implies  $\varphi(X_{+1}) \subset S_{-3}$ . Indeed, otherwise the intersection of  $\varphi_{+1}(X_{+1})$  with each of the connected components of the variety of  $\mathbb{P}^3$ -s on  $S$  will be closed, and consequently  $X_{+1}$  will be the disjoint union of two non-empty closed subvarieties, the preimages of those intersections. This is impossible, and hence  $\varphi_{+1}(X_{+1}) \subset S_{-3}$ . Moreover, the morphism  $\varphi_{+1} : X_{+1} \rightarrow S_{-3}$  is an embedding since distinct  $\mathbb{P}^3$ -s on  $X$  are mapped by  $\varphi$  to distinct  $\mathbb{P}^3$ -s on  $S$ .

Let  $\pi_S : S \rightarrow \mathbb{P}(V_S)$  be the minimal projective embedding. The composition  $\pi := \pi_S \circ \varphi$  fits into a commutative diagram of embeddings

$$\begin{array}{ccc} X & \xrightarrow{\varphi} & S \\ \pi \searrow & & \downarrow \pi_S \\ & & \mathbb{P}(V_S). \end{array}$$

Since distinct maximal  $\mathbb{P}^3$ -s on  $S$  determine distinct points in  $G(4, V_S)$ , the above diagram induces a commutative diagram of embeddings

$$(15) \quad \begin{array}{ccc} X_{+1} & \xrightarrow{\varphi_{+1}} & S_{-3} \\ & \searrow \pi^+ & \downarrow \pi_S^{-3} \\ & & G(4, V_S). \end{array}$$

We show now that the existence of such a diagram is contradictory.

Let us first observe that  $(\pi_S^{-3})^*(\mathcal{O}_{G(4, V_S)}(1)) \cong \mathcal{O}_{S_{-3}}(2)$ . To justify this it suffices to show that the pullback of  $\mathcal{O}_{G(4, V_S)}(1)$  to any maximal standard quadric  $Q$  on  $S_{-3}$  is isomorphic to  $\mathcal{O}_Q(2)$ . A maximal standard quadric on  $S_{-3}$  is 6-dimensional and has the form  $Q^6 = Q_{S_{-3}, W_{l-4}}^6$  for a unique  $W_{l-4} \in GO(l-4, V_{2l})$ . Fix  $W_{l-4}$  and let  $\tilde{S} = \{W \in S : W_{l-4} \subset W\}$ . Then  $\tilde{S}$  is isomorphic to the spinor Grassmannian  $GO(4, V_8)$ , where  $V_8 := W_{l-4}^\perp / W_{l-4}$ , and the inclusion  $\tilde{S} \subset S$  is a standard extension. The maximal  $\mathbb{P}^3$ -s on  $S$  arising from points in  $Q^6$  are all contained in  $\tilde{S}$ , and hence in  $\mathbb{P}(V_{\tilde{S}}) \subset \mathbb{P}(V_S)$ . We have  $\pi_S^{-3}(Q^6) \subset G(4, V_{\tilde{S}}) \subset G(4, V_S)$ . Thus without loss of generality we may assume  $l = 4$  and consider the minimal projective embedding  $\pi_{\tilde{S}} : \tilde{S} \rightarrow \mathbb{P}(V_{\tilde{S}})$  instead of  $\pi_S$ . Since  $\tilde{S} \cong GO(1, V_8) \cong Q^6$  the embedding  $\pi_{\tilde{S}}$  is isomorphic to the tautological embedding  $\tau : Q^6 \hookrightarrow \mathbb{P}(V_8)$ . There are two families of maximal  $\mathbb{P}^3$ -s on  $Q^6$ , parametrized respectively by  $\tilde{S} = \tilde{S}^+$  and  $\tilde{S}^-$ . The respective embeddings  $\tilde{S}^{\pm 1} \hookrightarrow G(4, V_8)$ , one of which replaces  $\pi_{\tilde{S}}^{-3}$  when  $Q^6$  replaces  $\tilde{S}$ , are the tautological embeddings which are well known to satisfy the claimed property  $(\tau^\pm)^*(\mathcal{O}_{G(4, V_8)}(1)) \cong \mathcal{O}_{\tilde{S}^\pm}(2)$  (see Subsection 1.1).

Next we check that  $(\pi^+)^*(\mathcal{O}_{G(4, V_S)}(1)) \cong \mathcal{O}_{X_{+1}}(3)$ . The projective embedding  $\pi_S \circ \varphi : X \rightarrow \mathbb{P}(V_S)$  factors through the minimal projective embedding  $\pi_X : X \hookrightarrow \mathbb{P}(V_X)$  which in turn induces an embedding  $\pi_X^+ : X_{+1} \hookrightarrow G(4, V_X)$ . We have  $(\pi^+)^*(\mathcal{O}_{G(4, V_S)}(1)) \cong (\pi_X^+)^*(\mathcal{O}_{G(4, V_X)}(1))$ . Suppose first that  $X$  is an ordinary Grassmannian  $G(3, V_n)$ , so that  $X_{+1} = G(4, V_n)$ . Then  $V_X = \Lambda^3 V_n$  and the map  $\pi_X^+$  is given by

$$\pi_X^+ : X_{+1} = G(4, V_n) \hookrightarrow G(4, \Lambda^3 V_n), \quad U \mapsto \Lambda^3 U.$$

The pullback  $(\pi_X^+)^*\mathcal{S}_4$  of the tautological bundle  $\mathcal{S}_4$  on  $G(4, \Lambda^3 V_n)$  is isomorphic to the exterior power  $\Lambda^3 \tilde{\mathcal{S}}_4$  of the tautological bundle  $\tilde{\mathcal{S}}_4$  on  $G(4, V_n)$ . Hence the determinant line bundle  $\Lambda^4(\pi_X^+)^*\mathcal{S}_4$  is isomorphic to  $\mathcal{O}_{X_{+1}}(-3)$ , and  $\Lambda^4(\pi_X^+)^*\mathcal{S}_4^*$  is isomorphic to  $\mathcal{O}_{X_{+1}}(3)$ . The cases when  $X$  is a regular orthogonal or symplectic Grassmannian are handled by considering an isotropic extension  $Y = G(3, V_{\lfloor n/2 \rfloor}) \xrightarrow{\iota} X$ , and reducing to the case of the ordinary Grassmannian  $Y$ .

Since the integers 2 and 3 are relatively prime we must conclude that the diagram (15) is contradictory. Hence, as asserted in the proposition, the morphism  $\varphi_+ : X_{+1} \rightarrow S^-$  is well defined also if  $X_{+1}$  parametrizes maximal projective spaces on  $X$  of dimension 3.

To construct the morphism  $\varphi_{-1} : X_{-1} \rightarrow S^-$  we need to consider cases. If  $X$  is an ordinary Grassmannian the definition of  $\varphi_{-1}$  is completely analogous to that of  $\varphi_{+1}$ , due to the isomorphism  $G(m, V_n)_{-1} \cong G(n-m, V_n^*)_{+1}$ . For a symplectic Grassmannian the situation is similar because  $X_{-1}$  parametrizes a family of maximal projective spaces on  $X$ , and if  $P$  belongs to  $X_{-1}$  then a morphism  $\varphi_{-1}$  is obtained analogously to  $\varphi_{+1}$ . If  $X$  is an orthogonal Grassmannian, then  $X_{-1}$  parametrizes the maximal standard quadrics on  $X$ . To establish that  $\varphi_{-1} : X_{-1} \rightarrow S^-$  is a well-defined morphism we have to show that for every  $U_{m-1} \in X_{-1}$  the maximal standard quadric  $Q = Q_{X, U_{m-1}}$  on  $X$  is mapped by  $\varphi$  into a projective space  $\mathbb{P}_{S, W_l}^{l-1}$  on  $S$ . Since  $X$  is regular we have  $\dim Q \geq 5$ . Lemma 3.5 implies that  $\varphi|_Q$  factors through  $\mathbb{P}_{S, W_l}^{l-1}$  as claimed, unless  $\dim Q$  is equal to 5 or 6 and  $\varphi(Q)$  is contained in  $Q_{S, W_{l-4}}^6$  for some  $W_{l-4} \in GO(l-4, V_{2l})$ .

We show now that the latter is impossible. Assume to the contrary that  $\varphi(Q) \subset Q_{S, W_{l-4}}^6$  for some  $W_{l-4} \in GO(l-4, V_{2l})$ . It suffices to consider the smallest case  $X = GO(2, V_9)$  and  $\dim Q = 5$ . Let  $U_2 \in X$ , and let  $U_1^{(1)}, U_1^{(2)} \in GO(1, V_9) = X_{-1}$  be distinct isotropic lines contained in  $U_2$ . Then  $\{U_2\} = Q_{X, U_1^{(1)}}^5 \cap Q_{X, U_1^{(2)}}^5$  and  $\varphi(Q_{X, U_1^{(i)}}^5) \subset Q_{S, W_{l-4}^{(i)}}^6$  where  $W_{l-4}^{(1)}, W_{l-4}^{(2)} \in$

$GO(l-4, V_{2l})$  are distinct, because otherwise the intersection of the two images would have positive dimension. Consider projective planes  $P_X^{(i)} \subset Q_{X, U_1^{(i)}}^5$  passing through  $U_2$ , for  $i = 1, 2$ . Let  $U_3^{(3)}, U_3^{(4)} \in GO(3, V_9) = X_{+1}$  be distinct 3-dimensional isotropic subspaces containing  $U_2$ . Then the two planes  $P_X^{(j)} := \mathbb{P}_{X, U_3^{(j)}}^2$  for  $j = 3, 4$  intersect at the point  $U_2$ . For  $i = 1, 2$  and  $j = 3, 4$  the intersection  $P_X^{(i)} \cap P_X^{(j)}$  is a projective line on  $X$  passing through  $U_2$ . Thus the images  $P^{(j)} := \varphi(P_X^{(j)})$  in  $S$  satisfy the hypothesis of Lemma 3.6 which implies that  $P^{(3)} \cup P^{(4)}$  is contained in a projective of the form  $\mathbb{P}_{S, \bar{W}_l}^{l-1}$  for  $\bar{W}_l \in S^-$  which is necessarily unique. It follows that  $\varphi_{+1}(U_3^{(3)}) = \varphi_{+1}(U_3^{(4)}) = \bar{W}_l = \varphi_{+1}(U_3)$  for all  $U_3 \in X_{+1}$ , because any two points  $U_3, U_3' \in X_{+1}$  are connected by a sequence of points  $U_3 = U_3^0, \dots, U_3^k = U_3'$  such that  $\mathbb{P}_{X, U_3^j}^2 \cap \mathbb{P}_{X, U_3^{j-1}}^2$  is a point in  $X$ . This means that the morphism  $\varphi_{+1}$  is constant and  $\varphi$  factors through the projective space  $\mathbb{P}_{S, \bar{W}_l}^{l-1}$ , which contradicts the hypothesis of the proposition. This contradiction implies that the morphism  $\varphi_{-1} : X_{-1} \rightarrow S^-$  is well defined for  $X = GO(2, V_9)$  and thus for all regular Grassmannians.

Let us remark for further use that the morphisms  $\varphi_{\pm 1}$  are defined under the weaker assumption that  $\varphi$  is a linear morphism because the restriction of any linear morphism  $X \rightarrow S$  to any projective space  $P$  on  $X$  is a linear embedding  $P \hookrightarrow S$ .

Our next task is to show that the morphisms  $\varphi_{\pm 1} : X_{\pm 1} \rightarrow S^-$  are linear embeddings. We concentrate on the case of  $\varphi_{-1}$  as the case of  $\varphi_{+1}$  is similar. Recall that a projective line on  $X$  corresponds to a unique flag  $U_{m-1} \subset U_{m+1}$  with  $U_{m-1} \in X_{-1}$  and  $U_{m+1} \in X_{+1}$ . Since  $\varphi$  is linear we have  $\varphi(\mathbb{P}_{X, U_{m-1} \subset U_{m+1}}^1) \subset \mathbb{P}_{S, \varphi_{+1}(U_{m+1})}^{l-1} \cap \mathbb{P}_{S, \varphi_{-1}(U_{m-1})}^{l-1}$ . Hence

$$(16) \quad \dim \varphi_{+1}(U_{m+1}) \cap \varphi_{-1}(U_{m-1}) = l - 2,$$

and the images  $\varphi_{-1}(U_{m-1})$  and  $\varphi_{+1}(U_{m+1})$  span a projective line on  $S^-$ .

The variety  $X_{-2, +1}$  of flags of subspaces  $U_{m-2} \subset U_{m+1}$  in  $V_n$  (of the indicated dimensions) parametrizes a family of projective planes on  $X$ , and a family of projective planes on  $X_{-1}$ . The subspaces  $U_{m-2}$  and  $U_{m+1}$  are assumed isotropic if  $X = GO(m, V_n)$  or  $X = GS(m, V_n)$ . Fix  $(U_{m-2} \subset U_{m+1}) \in X_{-2, +1}$ , and let  $P := \mathbb{P}_{X, U_{m-2} \subset U_{m+1}}^2$  and  $P_{-1} := \mathbb{P}_{X_{-1}, U_{m-2} \subset U_{m+1}}^2$  be the respective projective planes on  $X$  and  $X_{-1}$ . Set  $\bar{W}_l := \varphi_{+1}(U_{m+1})$  and let  $W_{l-3} \subset \bar{W}_l$  be such that

$$\varphi(\mathbb{P}_{X, U_{m-2} \subset U_{m+1}}^2) = \mathbb{P}_{S, W_{l-3} \subset \bar{W}_l}^2.$$

We aim to describe the image  $\varphi_{-1}(P_{-1})$ . Let  $U^1, U^2, U^3 \in P$  be three non-collinear points and let  $U^{ij} := U^i \cap U^j \in P_{-1}$  for  $1 \leq i < j \leq 3$ . Set  $W^i := \varphi(U^i)$  and  $\bar{W}^{ij} := \varphi_{-1}(U^{ij})$ . By (16),  $\bar{W}^{ij}$  and  $\bar{W}_l$  belong to a projective line on  $S^-$ , i.e.,  $\bar{W}^{ij} \cap \bar{W}_l$  has dimension  $l-2$ . On the other hand, the space  $U^{ij}$  corresponds to a projective line  $\mathbb{P}_{X, U^{ij} \subset U_{m+1}}^1 \subset P$  on  $X$  passing through  $U^i$  and  $U^j$ . The image under  $\varphi$  of  $\mathbb{P}_{X, U^{ij} \subset U_{m+1}}^1$  is the projective line on  $S$  passing through  $W^i$  and  $W^j$ , and hence

$$W_{l-3} \subset W^i \cap W^j = \bar{W}^{ij} \cap \bar{W}_l.$$

It follows in particular that  $\varphi_{-1}(P_{-1})$  is contained in the projective space  $\mathbb{P}_{S^-, W_{l-3}}^3$  on  $S^-$ . Therefore the points  $\bar{W}^{12}, \bar{W}^{13}, \bar{W}^{23} \in S^-$  are pairwise collinear, and the intersections  $W_{l-2}^1 := \bar{W}^{12} \cap \bar{W}^{13}$ ,  $W_{l-2}^2 := \bar{W}^{12} \cap \bar{W}^{23}$ ,  $W_{l-2}^3 := \bar{W}^{23} \cap \bar{W}^{13}$  are  $(l-2)$ -dimensional, defining the respective lines.

We claim that the triple  $\bar{W}^{12}, \bar{W}^{13}, \bar{W}^{23}$  is not collinear. Indeed, the  $(l-2)$ -dimensional spaces  $W_{l-2}^j$ ,  $j = 1, 2, 3$  also define projective lines on  $S$  contained in  $\mathbb{P}_{S, W_{l-3}}^3$ , and the points  $W^j$  are obtained as the intersection of these lines with  $\varphi(P)$ , i.e.,  $\{W^j\} = \mathbb{P}_{S, W_{l-2}^j}^1 \cap \varphi(P)$ . Since the triple  $W^1, W^2, W^3 \in \varphi(P)$  is not collinear, the three lines are distinct, and so the three spaces  $W_{l-2}^j$  are distinct. Therefore the points  $\bar{W}^{12}, \bar{W}^{13}, \bar{W}^{23} \in S^-$  are not collinear as

claimed, and are contained in a unique projective plane on  $S^-$  of the form  $\mathbb{P}_{S^-, W_{l-3} \subset W_l}^2$  for some  $W_l \in S$ . Moreover, the equality

$$W_l = W_{l-2}^1 + W_{l-2}^2 + W_{l-2}^3$$

holds. Indeed,  $W_l$  necessarily contains  $W_{l-2}^j$ , we have  $W_{l-2}^1 \cap W_{l-2}^2 \cap W_{l-2}^3 = W_{l-3}$ , and the three intersections  $W_l \cap \bar{W}^{ij} = W_{l-2}^i + W_{l-2}^j$  for  $1 \leq i < j \leq 3$  are distinct  $(l-1)$ -dimensional subspaces  $W_l$ , each contained in the respective  $\bar{W}^{ij}$ .

We show now that  $W_l$  depends only on  $P$  and not on the choice of the three points  $U^1, U^2, U^3$  in  $P$ . Indeed, it is enough to check that  $W_l$  remains unchanged if the triple  $U^1, U^2, U^3$  is replaced by  $U', U^2, U^3$  with  $U' \in \mathbb{P}_{X, U^{12} \subset U_{m+1}}^1 \setminus \{U^1, U^2\}$ . The corresponding triple in  $P_{-1}$  is  $U^{12} = U' \cap U^1 = U' \cap U^2, U'' := U' \cap U^3, U^{23}$ , and we have  $U'' \in \mathbb{P}_{X_{-1}, U_{m-2} \subset U^3}^1 \setminus \{U^{13}, U^{23}\}$ . The latter triple's image in  $S^-$  under  $\varphi_{-1}$  is the triple  $\bar{W}^{12}, \bar{W}'' := \varphi_{-1}(U''), \bar{W}^{23}$ . Going back to the space  $W_l$ , we observe that  $W_l$  is the sum of its intersections with any two of the spaces  $\bar{W}^{12}, \bar{W}^{13}, \bar{W}^{23}$ ; in particular  $W_l = W_l \cap \bar{W}^{12} + W_l \cap \bar{W}^{23}$ . Hence the triple  $\bar{W}^{12}, \bar{W}'', \bar{W}^{23}$  determines the same point  $W_l \in S$  (and hence the same plane in  $S^-$ ) as the triple  $\bar{W}^{12}, \bar{W}'', \bar{W}^{23}$ . It follows that  $\varphi_{-1}(P_{-1}) \subset \mathbb{P}_{S^-, W_{l-3} \subset W_l}^2$ . Moreover, since every projective line on  $\varphi(P)$  is necessarily the form  $\mathbb{P}_{S, \varphi_{-1}(U) \cap \bar{W}_l}^1$  for some  $U \in P_{-1}$ , we have an equality  $\varphi_{-1}(P_{-1}) = \mathbb{P}_{S^-, W_{l-3} \subset W_l}^2$ .

To prove that the morphism  $\varphi_{-1}$  is linear it suffices to prove that its restriction to a projective line on  $P_{-1}$  is linear. The projective lines on  $P_{-1}$  are parametrized by the points of  $P$ . The projective line  $\mathbb{P}_{X_{-1}, U_{m-2} \subset U^1}^1$  corresponding to  $U^1$  is spanned by  $U^{12}$  and  $U^{13}$ . Note that  $W^1 \cap W_l = W_{l-2}^1 = \bar{W}^{12} \cap \bar{W}^{13}$ . In particular the intersection  $\bar{W}^{12} \cap \bar{W}^{13}$  does not depend on the choice of the two points on  $\mathbb{P}_{X_{-1}, U_{m-2} \subset U^1}^1$ . Thus the image of this projective line under  $\varphi_{-1}$  is the line  $\mathbb{P}_{S^-, W_{l-2}^1}^1$ , and we conclude that  $\varphi_{-1}$  is a linear morphism as claimed.

It remains to show that  $\varphi_{-1}$  is an embedding. If  $m \geq 3$  then  $X_{-1}$  is a regular Grassmannian and  $X = (X_{-1})_{+1}$ , so  $\varphi$  can be recovered from  $\varphi_{-1}$  as  $\varphi = (\varphi_{-1})_{+1}$ , assuming only that  $\varphi_{-1}$  is a linear morphism. This implies the injectivity of  $\varphi_{-1}$  as follows. Any linear morphism is injective on projective spaces, so we assume  $U_{m-1}, U'_{m-1} \in X_{-1}$  are not collinear and  $\varphi_{-1}(U_{m-1}) = \varphi_{-1}(U'_{m-1}) =: \bar{W}_l \in S^-$ . Let  $U_m, U'_m \in X$  be such that  $U_{m-1} \subset U_m$  and  $U'_{m-1} \subset U'_m$ . The non-collinearity of  $U_{m-1}, U'_{m-1}$  implies  $U_m \neq U'_m$ . We have  $\bar{W}_l \in \varphi_{-1}(\mathbb{P}_{X_{-1}, U_m}^{m-1}) \cap \varphi_{-1}(\mathbb{P}_{X_{-1}, U'_m}^{m-1})$  and hence  $\varphi(U_m), \varphi(U'_m) \in \mathbb{P}_{S, \bar{W}_l}^{l-1}$ . In particular  $\varphi(U_m), \varphi(U'_m)$  are contained in a projective line  $L$  on  $S$ . Since  $\varphi$  is linear,  $L$  is the image of a projective line  $L'$  on  $X$  containing  $U_m, U'_m$ . In turn,  $L' \subset X$  is contained in a unique maximal projective space  $\mathbb{P}_{X, U}^d$  (or maximal standard quadric if  $X$  is an orthogonal Grassmannian) for a some  $U \in X_{-1}$ , which implies  $U = U_{m-1} = U'_{m-1}$ . Since an injective linear morphism is an embedding, we infer that  $\varphi_{-1}$  is an embedding for  $m \geq 3$ .

If  $m = 2$  and the Grassmannian  $X$  is ordinary or symplectic, then  $X_{-1}$  is a projective space and the linear morphism  $\varphi_{-1}$  is automatically an embedding. The remaining case is  $X = GO(2, V_n)$ , where  $X_{-1} = GO(1, V_n)$  is a quadric. Here  $X$  can be viewed as the parameter space for the family of projective planes on  $X_{-1}$ , and we can recover  $\varphi$  from  $\varphi_{-1}$  as the morphism mapping a projective plane  $P_{-1}$  on  $X_{-1}$  to the unique  $W \in S$  for which  $\varphi_{-1}(P_{-1}) \subset \mathbb{P}_{S^-, W}^{l-1}$ . An argument similar to the case  $m \geq 3$  implies that  $\varphi_{-1}$  is injective and hence an embedding.  $\square$

Next we describe the embeddings  $(\theta_l^k)_{\pm 1}$  for the embeddings  $\theta_l^k$  introduced in Subsection 3.1. In their definition,  $V_l$  was identified with a maximal isotropic subspace of  $V_{2l}$ , which belongs to  $S$  if  $k$  is even (see (6)), and to  $S^-$  if  $k$  is odd (see (7)). This was convenient to keep the target space  $S$  fixed for all  $k$ . Here it will be convenient to keep a maximal isotropic subspace  $V_l \subset V_{2l}$  fixed, and allow the target space of  $\theta_l^k$  to be  $S^+$  or  $S^-$  depending on the parity of  $k$ .

**Lemma 3.8.** *Let  $V_{2l} = V_l \oplus V_l^*$  be a splitting of an orthogonal space  $V_{2l}$  into a sum of isotropic subspaces. The embeddings*

$$\theta_l^k : G(l - k, V_l) \rightarrow S^{\text{sign}(-1)^k}, \quad U \rightarrow U \oplus U^\perp \cap V_l^*$$

*satisfy  $(\theta_l^k)_{\pm 1} = \theta_l^{k \mp 1}$ .*

*Proof.* The definition of the morphisms  $\theta_l^k$  implies

$$\theta_l^k(\mathbb{P}_{X, U_{l-k\pm 1}}) \subset \mathbb{P}_{S, \theta_l^{k\pm 1}(U_{l-k\pm 1})},$$

and the statement follows immediately.  $\square$

**Proposition 3.9.** *Let  $X \xrightarrow{\varphi} S$  be a linear embedding of a regular Grassmannian  $X$  into  $S$ . Then  $\varphi$  factors through one of the embeddings  $\theta_l^k : G(l - k, V_l) \hookrightarrow S$  for a suitable isotropic subspace  $V_l \subset V_{2l}$  and  $1 \leq k \leq l - 2$ .*

*Proof.* The proof proceeds by induction on  $m$ . For  $m = 2$  the Grassmannian  $X_{-1}$  is a projective space or a quadric, and the embedding  $\varphi_{-1} : X_{-1} \rightarrow S^-$  factors through a unique projective space  $P := \mathbb{P}_{S^-, W_l}^{l-1}$  for some  $W_l \in S$ . The variety  $X$  can be interpreted as the parameter space for a family of projective lines on  $X_{-1}$ . Hence  $\varphi(X) \subset \{W \in S : \dim W \cap W_l = l - 2\}$ . We will now show that  $\varphi$  factors through an embedding  $\theta_l^2$ . For this it suffices to find an isotropic subspace  $W_l^* \subset V_{2l}$  such that  $\dim \varphi(U) \cap W_l^* = 2$  for all  $U \in X$ .

Claim: if  $W_l^* \subset V_{2l}$  is an isotropic subspace such that  $V_{2l} = W_l \oplus W_l^*$ , then  $\dim \bar{W} \cap W_l^* = 1$  for all  $\bar{W} \in P$ . Recall that the isotropic subspaces of  $V_{2l}$  complementary to  $W_l$  all belong to  $S^-$  if  $l$  is even, and all belong to  $S$  if  $l$  is odd. Consequently, for any  $\bar{W} \in S^-$  the dimension of the intersection  $\bar{W} \cap W_l^*$  is odd. On the other hand, since  $\bar{W} \cap W_l$  has dimension  $l - 1$ , the space  $\bar{W} \cap W_l^*$  has dimension at most 1. Thus  $\dim \bar{W} \cap W_l^* = 1$ .

It follows that every  $W_l^*$  as above satisfies  $\dim \varphi(U) \cap W_l^* = 2$  for all  $U \in X$ , which allows us to conclude that  $\varphi$  factors through the embedding  $\theta_l^2 : G(l - 2, W_l) \hookrightarrow S$  defined in (6) by the splitting  $V_{2l} = W_l \oplus W_l^*$ . The case  $m = 2$  is settled.

Now suppose that  $m > 2$  and assume that the statement holds for  $p < m$ . Then  $X_{-1}$  is a regular Grassmannian and the embedding  $\varphi_{-1} : X_{-1} \hookrightarrow S^-$  factors through some  $\theta_l^{k'}$ . Since  $(\varphi_{-1})_{+1} = \varphi$ , Lemma 3.8 implies that  $\varphi$  factors through  $\theta_l^k$  with  $k = k' + 1$  or  $k = k' - 1$ .  $\square$

We are now ready for the main result of this subsection.

**Theorem 3.10.** *Let  $S = GO(l, V_{2l})$  for  $l \geq 5$ , let  $X$  be a regular Grassmannian, a quadric, or a spinor Grassmannian, and let  $X \xrightarrow{\varphi} S$  be a linear embedding which does not factor through a projective space. Then all possible options are as follows:*

- (i)  $X$  is a spinor Grassmannian  $GO(m, V_{2m})$  with  $4 \leq m \leq l$  and  $\varphi$  is a standard extension.
- (ii)  $X = GO(1, V_7) \cong Q^5$  and  $\varphi$  is a composition

$$\varphi : X \cong Q^5 \xrightarrow{\sigma_1} Q^6 \cong GO(4, V_8) \xrightarrow{\sigma_2} GO(l, V_{2l})$$

where  $\sigma_1, \sigma_2$  are standard extensions.

- (iii)  $\varphi$  factors through one of the embeddings  $\theta_l^k$ ,  $2 \leq k \leq l - 2$ , defined in (6) and (7); in such a case exactly one of the following occurs:

- (iii.1)  $X \cong G(m, V_n)$  with  $2 \leq m \leq n/2$  and  $\varphi$  is a composition

$$\varphi : G(m, V_n) \xrightarrow{\sigma} G(l - k, V_l) \xrightarrow{\theta_l^k} GO(l, V_{2l})$$

where  $\sigma$  is a standard extension.

- (iii.2)  $X \cong GS(m, V_n)$  with  $1 < m < n/2$  and  $\varphi$  is a composition

$$\varphi : X \cong GS(m, V_n) \xrightarrow{\tau} G(m, V_n) \xrightarrow{\sigma} G(l - k, V_l) \xrightarrow{\theta_l^k} GO(l, V_{2l})$$

where  $\tau$  is a tautological embedding and  $\sigma$  is a standard extension.

(iii.3)  $X \cong GO(m, V_n)$  with  $1 < m < \lfloor \frac{n-3}{2} \rfloor$  and  $\varphi$  is a composition

$$\varphi : X \cong GO(m, V_n) \xrightarrow{\tau} G(m, V_n) \xrightarrow{\sigma} G(l-k, V_l) \xrightarrow{\theta_l^k} GO(l, V_{2l})$$

where  $\tau$  is a tautological embedding and  $\sigma$  is a standard extension.

*Proof.* The case of a spinor Grassmannian  $X = GO(m, V_{2m})$  is considered in Theorem 2.4, and the quadrics of dimension at least 5 are considered in Lemma 3.5.

It remains to address the case of a regular Grassmannian  $X$ . By Proposition 3.9,  $\varphi$  factors through one of the embeddings  $\theta_l^k$  and hence option (iii) holds. Now  $\varphi$  can be written as  $\varphi = \theta_l^k \circ \psi$  for a linear embedding  $\psi : X \hookrightarrow G(k, V_l)$ . The linear embeddings of regular Grassmannians into ordinary Grassmannians are classified in Theorems 2.4 and 2.5, and the resulting possibilities are the ones listed in (iii.1), (iii.2), (iii.3).  $\square$

**Remark 3.1.** If  $X = GS(m, V_{2m})$  for  $m \geq 2$ , or  $X = GO(m-2, V_{2m-\varepsilon})$  for  $5 \leq m$ ,  $\varepsilon \in \{0, 1\}$ , there are linear embeddings  $X \hookrightarrow S$  which factor through the tautological embedding  $\tau : X \hookrightarrow G(m, V_n)$  and have the form given in (iii.2) and (iii.3), respectively. For  $X = GO(m-2, V_{2m-\varepsilon})$  there are also linear embeddings of the form

$$\varphi : X \cong GO(m-2, V_{2m-\varepsilon}) \xrightarrow{\delta_m^\varepsilon} G(2, V_{2m-1}) \xrightarrow{\sigma} G(l-k, V_l) \xrightarrow{\theta_l^k} GO(l, V_{2l})$$

where  $\delta_m^0 = \delta_m$  is the embedding defined in (8),  $\delta_m^1$  is the embedding defined in (11), and  $\sigma$  is a standard extension. To complete the classification of linear embeddings of Grassmannians one needs to determine if other embeddings of  $X$  into  $S$  exist.

**3.5. Maximal linear embeddings of Grassmannians.** A linear embedding of Grassmannians  $X \xrightarrow{\varphi} Y$  is **maximal** if  $\varphi$  is a proper embedding, and does not factor through a Grassmannian  $Z$  properly contained in  $Y$  and properly containing  $\varphi(X)$ .

**Corollary 3.11.** Let  $X \xrightarrow{\varphi} Y$  be a maximal linear embedding of Grassmannians. Assume that  $Y$  is not a projective space or a quadric, and if  $Y$  is a spinor Grassmannian then  $X \neq GS(m, V_{2m})$  for  $m \geq 3$ ,  $X \neq GO(m-2, V_{2m-\varepsilon})$  for  $m \geq 4$ ,  $\varepsilon \in \{0, 1\}$ . Then, up to compositions with isomorphisms of  $X$  and  $Y$ ,  $\varphi$  is one of the following.

- (a) For  $Y = G(l, V_s)$  with  $1 < l \leq s/2$ :
  - (a.1) a standard extension  $G(m, V_{s-1}) \hookrightarrow G(l, V_s)$  with  $m \in \{l, l-1\}$ ;
  - (a.2) a tautological embedding  $GO(l, V_s) \hookrightarrow G(l, V_s)$  with  $s \neq 2l, 2l+1, 2l+2$ , or  $GS(l, V_s) \hookrightarrow G(l, V_s)$ .
- (b) For  $Y = GO(l, V_s)$  with  $1 < l < \lfloor \frac{s-1}{2} \rfloor$ :
  - (b.1) a standard extension  $GO(l, V_{s-1}) \hookrightarrow GO(l, V_s)$ , or  $GO(l-1, V_{s-2}) \hookrightarrow GO(l, V_s)$ ;
  - (b.2) an isotropic extension  $G(l, V_{\frac{s}{2}}) \hookrightarrow GO(l, V_s)$  where  $s$  is even and  $V_{\frac{s}{2}} \subset V_s$  is a maximal isotropic subspace;
  - (b.3) if  $l = 2$  and  $s = 2^{n-1}$  for some  $n \geq 1$ , the embedding  $\delta_n^O : GO(n-2, V_{2n}) \hookrightarrow GO(2, V_{2^{n-1}})$  defined in Subsection 3.2, whenever the minimal projective embedding of  $GO(n, V_{2n})$  factors through a quadric.
- (c) For  $Y = GS(l, V_{2s})$  with  $1 < l < s$ :
  - (c.1) an isotropic extension  $G(l, V_s) \hookrightarrow GS(l, V_{2s})$  where  $V_s \subset V_{2s}$  is a maximal isotropic subspace;
  - (c.2) a standard extension  $GS(m, V_{2s-2}) \hookrightarrow GS(l, V_{2s})$  with  $m \in \{l-2, l-1, l\}$ ;
  - (c.3) if  $l = 2$  and  $s = 2^{n-1}$  for some  $n \geq 2$ , the embedding  $\delta_n^S : GO(n-2, V_{2n}) \hookrightarrow GS(2, V_{2^{n-1}})$  given in (10) provided  $V_{2^{n-1}}$  admits a non-degenerate skew-symmetric bilinear form for which every projective line on  $GO(n, V_{2n}) \subset \mathbb{P}(V_{2^{n-1}})$  is isotropic.
- (d) A standard extension  $GS(l-1, V_{2l-2}) \hookrightarrow GS(l, V_{2l})$ .
- (e) For  $Y = GO(l, V_{2l})$  with  $l \geq 5$ :

- (e.1) a standard extension  $GO(l-1, V_{2l-2}) \hookrightarrow GO(l, V_{2l})$ ;
- (e.2) for  $1 \leq k \leq l-1$  the embeddings  $\theta_l^k : G(l-k, V_l) \hookrightarrow GO(l, V_{2l})$  defined in (6) and (7).

*Proof.* The list is derived using the classification of linear embeddings from Theorems 2.4, 2.5, 3.3, 3.4, Proposition 3.1, and Theorem 3.10.  $\square$

#### 4. EQUIVARIANCE OF LINEAR EMBEDDINGS

A (generalized) flag variety is a coset space of the form  $G/P$  where  $G$  is a connected semisimple complex algebraic group and  $P$  is a parabolic subgroup. The Picard group of  $G/P$  is isomorphic to  $\mathbb{Z}$  if and only if  $P$  is a maximal parabolic subgroup of  $G$ . For a Grassmannian  $X$  we denote by  $G_X$  the simply connected cover of the connected component of the identity of the automorphism group of  $X$ . Concretely,  $G_X = SL(V_n)$  for  $X = G(m, V_n)$ ;  $G_X = Spin(V_n)$  for  $X = GO(m, V_n)$  with  $n \neq 2m+1$ ;  $G_X = Sp(V_n)$  for  $X = GS(m, V_n)$  with  $n$  even and  $m \neq 1$ . Then, in all cases we have  $X = G_X/P_m$  where  $P_m$  is the subgroup of  $G_X$  fixing a point in  $X$ .

The classification of linear embeddings of Grassmannians given Sections 2 and 3 has the remarkable consequence that almost every linear embedding of Grassmannians is equivariant for an appropriate classical group. There are exceptions: embeddings which factor through standard quadrics or standard symplectic projective spaces, as well as certain embeddings of  $GO(n-2, V_{2n})$  and  $GO(n-1, V_{2n+1})$  into orthogonal or symplectic Grassmannians. We will show that these are the only exceptions.

**Definition 4.1.** *An embedding of Grassmannians  $X \xrightarrow{\varphi} Y$  is  $G$ -equivariant (or simply equivariant) if  $G$  is a subgroup of  $G_X$  acting transitively on  $X$  and there is a homomorphism  $f : G \rightarrow G_Y$  with finite kernel such that  $\varphi(gx) = f(g)\varphi(x)$  for  $g \in G$  and  $x \in X$ .*

For any Grassmannian  $X$  the minimal projective embedding  $\pi_X : X \rightarrow \mathbb{P}(V_X)$  is equivariant under  $G_X$ . Furthermore, note that the image  $\varphi(X) \subset Y$  of a  $G$ -equivariant embedding is a closed orbit of the subgroup  $f(G) \subset G_Y$ .

**Theorem 4.1.** *Let  $X$  and  $Y$  be Grassmannians such that  $X \not\cong GO(n-2, V_{2n-1}), GO(n-2, V_{2n})$  for  $n \geq 4$ , and additionally  $X \not\cong GS(n, V_{2n})$  for  $n \geq 3$  if  $Y$  is a spinor Grassmannian. Then every linear embedding  $X \xrightarrow{\varphi} Y$ , which does not factor through a projective space or a quadric, is  $G_X$ -equivariant.*

Instead of proving Theorem 4.1 directly, we will describe the images of embeddings as in Theorem 4.1 as orbits of suitable groups, and this will imply the claimed equivariance. First, we introduce the following short-hand terminology. Let  $G$  be a connected semisimple linear algebraic group. We say that a subgroup  $L \subset G$  is an  **$\mathcal{L}$ -subgroup** ( $\mathcal{L}$  for Levi) if it is a maximal semisimple connected subgroup of a parabolic subgroup of  $G$ . If an  $\mathcal{L}$ -subgroup  $L \subset G$  is infinitesimally simple, we say that  $L$  is an  **$\mathcal{LS}$ -subgroup**. An algorithm for description of the  $\mathcal{L}$ -subgroups of a given simple group is provided by Dynkin in [3].

**Remark 4.1.** *If follows from standard properties of parabolic subgroups that, if  $G_1 \xrightarrow{f_1} G_2 \xrightarrow{f_2} G_3$  are homomorphisms of semisimple complex algebraic groups such that  $f_1(G_1) \subset G_2$  and  $f_2(G_2) \subset G_3$  are  $\mathcal{L}$ -subgroups (respectively,  $\mathcal{LS}$ -subgroups) then the image  $f_2 \circ f_1(G_1) \subset G_3$  is also an  $\mathcal{L}$ -subgroup (respectively,  $\mathcal{LS}$ -subgroup).*

**Proposition 4.2.**

- (i) *Let  $X \xrightarrow{\varphi} Y$  be a linear embedding of Grassmannians as in Theorem 4.1, in particular  $\varphi$  does not factor through a projective space or a quadric. Let  $Z \subset Y$  be a closed orbit of an  $\mathcal{LS}$ -subgroup of  $G_Y$ , containing  $\varphi(X)$  and minimal with this property. Then  $\varphi(X) = Z$  or the embedding  $\varphi(X) \hookrightarrow Z$  is one of the following:*

- (i.1) a tautological embedding  $GO(m, V_n) \hookrightarrow G(m, V_n)$  or  $GS(m, V_n) \hookrightarrow G(m, V_n)$ , where  $\varphi$  is one of the following: an embedding of  $X$  into an ordinary Grassmannian; a (mixed) combination of standard and isotropic extensions;  $Y = GO(l, V_{2l})$  and  $\varphi$  is one of the embeddings given in Theorem 3.10, (iii.2) and (iii.3).
- (i.2) a standard extension  $GO(m, V_n) \hookrightarrow GO(l, V_{n+1})$  with  $l \in \{m, m+1\}$ , where  $\varphi$  is a standard extension of orthogonal non-spinor Grassmannians  $GO(m, V_n) \hookrightarrow GO(l, V_s)$  with  $s - n \equiv 1 \pmod{2}$ .
- (i.3) a standard extension  $Q^5 \xrightarrow{\sigma_1} Q^6$ , where  $\varphi$  is an embedding of the form  $X = Q^5 \xrightarrow{\sigma_1} Q^6 \cong GO(4, V_8) \xrightarrow{\sigma_2} GO(l, V_{2l}) = Y$  for standard extensions  $\sigma_1, \sigma_2$ .
- (ii) Let  $Y$  be a Grassmannian. Every non-constant closed orbit  $X \subset Y$  of an  $\mathcal{LS}$ -subgroup  $L \subset G_Y$  is a Grassmannian, and  $X$  is not contained in a projective space or a standard quadric on  $Y$ , unless  $X$  is itself a projective space or a standard quadric on  $Y$ . Furthermore, except in the case where  $Y = GS(m, V_{2m})$  for some  $m$  and  $L$  is isomorphic to  $SL(k)$  for some  $k \geq 2$ , the embedding  $\varphi$  of  $X$  into  $Y$  is linear and there are three options:
  - (ii.1)  $\varphi$  is a standard extension;
  - (ii.2)  $X$  is an ordinary Grassmannian,  $Y$  is a symplectic or non-spinor orthogonal Grassmannian, and  $\varphi$  factors as

$$X \cong G(m, V_n) \xrightarrow{\sigma} G(l, V_r) \xrightarrow{\psi} Y$$

where  $\sigma$  is a standard extension and  $\psi$  is an isotropic extension.

- (ii.3)  $X$  is an ordinary Grassmannian,  $Y = GO(l, V_{2l})$ , and  $\varphi$  factors as

$$X \cong G(m, V_n) \xrightarrow{\sigma} G(k, V_l) \xrightarrow{\theta_l^k} Y$$

for some  $1 \leq k \leq l-1$ , where  $\sigma$  is a standard extension and  $\theta_l^k$  is the embedding defined in (6), (7).

*Proof.* (i). We begin with the case where  $Y = G(l, V_s)$  and hence  $G_Y = SL(V_s)$ . If  $X = G(m, V_n)$  then  $\varphi$  is a standard extension, since we have assumed that  $\varphi$  does not factor through a projective space. The image of every standard extension is also the image of a strict standard extension, so we may assume that  $\varphi$  is a strict standard extension. Thus we have a decomposition  $V_s = V_n \oplus V'$  and  $\varphi$  is given by (2). This embedding is clearly  $SL(V_n)$ -equivariant for the homomorphism  $f_{n,s} : SL(V_n) \hookrightarrow SL(V_s)$  whose image consists of the elements preserving the decomposition  $V_s = V_n \oplus V'$  and acting by identity on  $V'$ . Moreover, this image is an  $\mathcal{LS}$ -subgroup of  $SL(V_s)$ , because it is a Levi component of a parabolic subgroup of  $SL(V_s)$  fixing a flag of the form  $V_n \subset V_n \oplus U_1 \subset \dots \subset V_n \oplus U_{s-n+1} \subset V_s$  in  $V_s$  where  $U_1 \subset \dots \subset U_{s-n-1} \subset V'$  is a maximal flag in  $V'$ . (The choice of the maximal flag  $U_1 \subset \dots \subset U_{s-n-1} \subset V'$  does not matter.) This completes the argument for embeddings of ordinary Grassmannians.

If  $X = GS(m, V_n)$  then by Theorem 2.5 the embedding  $\varphi$  factors through the tautological embedding  $\tau_X$  as

$$GS(m, V_n) \xrightarrow{\tau_X} G(m, V_n) \xrightarrow{\sigma} G(l, V_s),$$

where  $\sigma$  is a standard extension. By the above,  $\sigma(G(m, V_n))$  is a closed orbit  $Z$  of an  $\mathcal{LS}$ -subgroup of  $SL(V_s)$ , and the embedding  $\varphi(X) \hookrightarrow Z$  is as claimed in (i.1). The case where  $X = GO(m, V_n)$  with  $m < \frac{n}{2} - 2$  is analogous.

Next we assume that  $Y = GO(l, V_s)$  with  $l < \lfloor \frac{s-1}{2} \rfloor$ . Then  $G_Y = Spin(V_s)$  but the  $G_Y$ -action on both  $Y$  and  $\mathcal{O}_Y(1)$  actually reduces to  $SO(V_s)$ . If  $X = G(m, V_n)$  then by Theorem 2.5 the embedding  $\varphi$  factors as a composition  $\iota \circ \sigma$  of a standard extension  $\sigma$  of ordinary Grassmannians and an isotropic extension  $\iota$ . By Remark 4.1 and the already established results on ordinary Grassmannians, it suffices to consider the case of an isotropic extension, i.e.,  $\varphi = \iota$ . Let  $U_n \subset V_s$  be an isotropic subspace with a fixed isomorphism  $V_n \cong U_n$  defining  $\varphi$ . Then  $\varphi$  is

$SL(V_n)$ -equivariant for the homomorphism  $h_{n,s}^O : SL(V_n) \rightarrow SO(V_s)$  whose image consists of the elements preserving a decomposition  $V_s = V_r \oplus U_n \oplus U_n^*$  and acting by the identity on  $V_r$  and on  $\Lambda^n U_n$ , where  $U_n^* \subset V_s$  is an isotropic subspace dual to  $U_n$  and  $V_r$  is an orthogonal complement to  $U_n \oplus U_n^*$ . The image of  $h_{n,s}^O$  is an  $\mathcal{LS}$ -subgroup acting transitively on the image of  $\varphi$ .

If  $X = GO(m, V_n)$  then the embedding  $\varphi$  is either a standard extension or a combination of standard and isotropic extensions. We consider the two cases. If  $\varphi$  is a standard extension then there is an orthogonal decomposition  $V_s \cong V_n \oplus V_{s-n}$  such that  $\varphi$  is given by (2), and  $\varphi$  is  $SO(V_n)$ -equivariant for the homomorphism

$$(17) \quad f_{n,s}^O : SO(V_n) \rightarrow SO(V_s)$$

whose image consists of the elements preserving the decomposition  $V_s \cong V_n \oplus V_{s-n}$  and acting trivially on  $V_{s-n}$ . The image of  $f_{n,s}^O$  is an  $\mathcal{LS}$ -subgroup of  $G_Y$  if and only if  $s - n \equiv 0 \pmod{2}$ .

If  $s - n \not\equiv 1 \pmod{2}$  then we observe that  $\varphi$  factors as a composition of two standard extensions of the form

$$GO(m, V_n) \xrightarrow{\sigma_1} GO(m, V_{n+1}) \xrightarrow{\sigma_2} GO(l, V_s).$$

Thus the image  $Z := \sigma_2(GO(m, V_{n+1}))$  is the orbit of an  $\mathcal{LS}$ -subgroup of  $G_Y$ , and the inclusion  $\varphi(X) \subset Z$  is a standard extension as claimed in (i.2).

Now suppose that the embedding  $\varphi$  is a combination of standard and isotropic extensions

$$X = GO(m, V_n) \xrightarrow{\tau_X} G(m, V_n) \xrightarrow{\sigma} G(l, V_r) \xrightarrow{\iota} GO(l, V_s) = Y.$$

By the previous cases, the image  $Z := \iota \circ \sigma(G(m, V_n)) \subset Y$  is the orbit of an  $\mathcal{LS}$ -subgroup of  $SO(V_s)$  isomorphic to  $SL(V_n)$ . The inclusion  $\varphi(X) \subset Z$  is given by the tautological embedding of  $X$  as claimed in (i.1).

To complete the argument for the case where  $Y = GO(l, V_s)$  is a non-spinor orthogonal Grassmannian, it remains to consider the case  $X = GS(m, V_r)$ . Here Theorem 2.5 implies that the embedding  $\varphi$  is a mixed combination of standard and isotropic extensions, and the statement of (i.1) applies for the same reasons as for the above case of a combination of standard and isotropic extensions.

The case of a spinor Grassmannian  $Y = GO(l, V_{2l})$  is analogous the case of non-spinor orthogonal Grassmannians, the only difference being that the embeddings  $\theta_l^k : G(l-k, V_l) \hookrightarrow Y$  replace the isotropic embeddings of ordinary Grassmannians into orthogonal Grassmannians.

The case  $Y = GS(l, V_{2s})$  is analogous to the case of non-spinor orthogonal Grassmannians  $GO(l, V_s)$ , and we leave it to the reader.

For part (ii) we use the following alternative characterization of  $\mathcal{L}$ -subgroups, see [4, § 30.2]. Let  $G$  be a connected semisimple linear algebraic group. A subgroup  $L \subset G$  is an  $\mathcal{L}$ -subgroup if and only if  $L$  is equal to the derived subgroup  $(G_\gamma)'$  of the centralizer subgroup  $G_\gamma \subset G$  of a one-parameter subgroup  $\gamma : \mathbb{C}^\times \rightarrow G$ . If  $L \subset G$  is an  $\mathcal{L}$ -subgroup then the normalizer  $N_G(L)$  contains a maximal torus  $T$  of  $G$ . Now let  $Y = G/P$  be a flag variety of  $G$  and  $L \subset G$  be an  $\mathcal{L}$ -subgroup. An  $L$ -orbit  $L \cdot x \subset Y$  is closed if and only if  $x$  is a  $T$ -fixed point for some maximal torus  $T \subset G$  contained in the normalizer  $N_G(L)$ . For any maximal torus  $T \subset G$ , the set of  $T$ -fixed points  $Y^T$  forms an orbit of the Weyl group  $W := N_G(T)/T$ .

We consider first the case  $Y = GS(l, V_{2s})$  where  $G_X = Sp(V_{2s})$ . Every  $\mathcal{LS}$ -subgroup of  $Sp(V_{2s})$  is isomorphic to  $SL(n)$  or  $Sp(2n)$  for some  $n \leq s$ , and is obtained via the following construction. Fix a decomposition of  $V_{2s}$  as a sum of two maximal isotropic subspaces,  $V_{2s} = U_s \oplus U_s^*$ . Let  $V_{2s} = V_{2n} \oplus V'$  be an orthogonal decomposition such that  $U_n := V_{2n} \cap U_s$  and  $U_n^* := V_{2n} \cap U_s^*$  are complementary maximal isotropic spaces of  $V_{2n}$ . We denote by  $L_{n,A}$  the subgroup of  $Sp(V_{2s})$  which preserves the decomposition  $V_{2s} = U_n \oplus U_n^* \oplus V'$  and acts by the identity on both  $V'$  and  $\Lambda^n U_n$ , and by  $L_{n,C}$  the subgroup of  $G$  preserving the decomposition

$V_{2s} = V_{2n} \oplus V'$  and acting by the identity on  $V'$ . Then there are isomorphisms

$$h_{n,2s}^S : SL(U_n) \cong L_{n,A} \subset Sp(V_{2s}) \quad \text{and} \quad f_{2n,2s}^S : Sp(V_{2n}) \cong L_{n,C} \subset Sp(V_{2s}) .$$

Next we observe that the closed orbits in  $Y$  of an  $\mathcal{LS}$ -subgroup  $L_{n,A} \subset Sp(V_{2s})$  are exactly the  $L_{n,A}$ -orbits of points  $U \in GS(l, V_{2s})$  such that  $U = (U \cap U_n) \oplus (U \cap U_n^*) \oplus (U \cap V')$ . Let us fix one such  $U$  and denote the closed orbit  $L_{n,A} \cdot U$  by  $X$ . Set  $m := \dim U \cap U_n$  and  $m^* := \dim U \cap U_n^*$ . Then  $X$  is an  $L_{n,A}$ -fixed point if and only if at least one of  $m$  and  $m^*$  is equal to 0. Suppose that  $X$  is not a point and  $m > 0$ , the case  $m^* > 0$  being analogous. Then  $X$  is isomorphic to the Grassmannian  $G(m, U_n)$ , since  $X$  is the image of the  $SL(U_n)$ -equivariant embedding

$$G(m, U_n) \xrightarrow{\varphi} GS(l, V_{2s}) , \quad \varphi(g \cdot (U \cap U_n)) := h_{n,2s}^S(g) \cdot U \quad \text{for } g \in SL(U_n) .$$

It is straightforward to check that the embedding  $\varphi$  is linear if and only if  $l \neq s$ . Furthermore, if  $\varphi$  is linear then it is a composition of a standard extension  $G(m, U_n) \hookrightarrow G(l, U_s)$  and an isotropic extension  $G(l, U_s) \hookrightarrow GS(l, V_{2s})$ , as claimed in (ii.2).

For an  $\mathcal{LS}$ -subgroup  $L_{n,C} \subset Sp(V_{2n})$ , the closed  $L_{n,C}$ -orbits in  $Y$  are exactly the  $L_{n,C}$ -orbits of points  $U \in GS(l, V_{2s})$  such that  $U = (U \cap V_{2n}) \oplus (U \cap V')$ . Let  $X := L_{n,C} \cdot U$  be a closed orbit and let  $m := \dim(U \cap V_{2n})$ . Then  $X$  is a fixed point if and only if  $m = 0$ . Otherwise,  $X$  is isomorphic to the Grassmannian  $GS(m, V_{2n})$  and is linearly embedded in  $Y$  by the  $Sp(V_{2n})$ -equivariant embedding

$$GS(m, V_{2n}) \xrightarrow{\varphi} GS(l, V_{2s}) , \quad \varphi(g \cdot (U \cap V_{2n})) := f_{2n,2s}^S(g) \cdot U \quad \text{for } g \in Sp(V_{2n})$$

which is a standard extension. This completes the argument for a symplectic Grassmannian  $Y$ .

The other cases are similar, and here we present the details only for a spinor Grassmannian. Suppose that  $Y = GO(l, V_{2l})$  with  $l \geq 5$ . Here  $G_Y = Spin(V_{2l})$  and the  $G_Y$ -action on  $\mathcal{O}_Y(1)$  does not reduce to an  $SO(V_{2l})$ -action. Every proper  $\mathcal{LS}$ -subgroup  $L \subset G_Y$  is isomorphic either to  $SL(n)$  for  $2 \leq n \leq l$ , or to  $Spin(2n)$  for  $3 \leq n < l$ . The inclusion of  $L$  in  $G$  is respectively

$$h_{n,2s}^O : SL(U_n) \cong L_{n,A} \subset G_Y \quad \text{or} \quad f_{2n,2l}^O : Spin(V_{2n}) \cong L_{n,D} \subset G_Y ,$$

where the subgroup  $L_{n,D}$  consists of the elements of  $Spin(V_{2l})$  preserving a fixed orthogonal decomposition  $V_{2l} = V_{2n} \oplus V_{2l-2n}$  and acting by the identity on  $V_{2l-2n}$ , and the subgroup  $L_{n,A} \subset Spin(V_{2l})$  consists of the elements preserving a fixed decomposition  $V_{2l} = U_n \oplus U_n^* \oplus V_{2l-2n}$  with  $U_n, U_n^* \subset V_{2l}$  isotropic and  $V_{2l-2n} = U_n^\perp \oplus (U_n^*)^\perp$ , and acting by the identity on both  $V_{2l-2n}$  and  $\Lambda^n U_n$ .

The closed  $L_{n,A}$ -orbits in  $Y = GO(l, V_{2l})$  are exactly the  $L_{n,A}$ -orbits of points  $U \in Y$  such that  $U = (U \cap U_n) \oplus (U \cap U_n^*) \oplus (U \cap V_{2l-2n})$ . Let us fix one such  $U$  and denote by  $X := L_{n,A} \cdot U \subset Y$  the closed  $L_{n,A}$ -orbit. Set  $m := \dim U \cap U_n$  and  $m^* := \dim U \cap U_n^*$ . Then  $X$  is an  $L_{n,A}$ -fixed point if and only if  $(m, m^*) \in \{(0, 0), (l, 0), (0, l)\}$ . Suppose that  $X$  is not a point and  $m > 0$  (the case  $m^* > 0$  is analogous). Then  $X$  is isomorphic to the Grassmannian  $G(m, U_n) \cong SL(U_n)/P_m$ , since  $X$  is the image of the  $SL(U_n)$ -equivariant embedding

$$G(m, U_n) \xrightarrow{\varphi} GO(l, V_{2l}) , \quad \varphi(g \cdot (U \cap U_n)) := f_{n,2l}^O(g) \cdot U \quad \text{for } g \in SL(U_n) .$$

The map  $\varphi$  is a linear embedding because it factors as a composition

$$X = G(m, U_n) \xrightarrow{\sigma} G(k, U_l) \xrightarrow{\theta_l^{l-k}} GO(l, V_{2l}) = Y$$

for some maximal isotropic subspace  $U_l \subset V_{2l}$  containing  $U_n$ , where  $\sigma$  is a standard extension and  $\theta_l^{l-k}$  is one of the embeddings defined in (6) and (7). Thus statement (ii.3) applies for the closed  $L_{n,A}$ -orbits in  $Y$ .

The closed  $L_{n,D}$ -orbits in  $Y = GO(l, V_{2l})$  are exactly the  $L_{n,D}$ -orbits through points  $U \in Y$  such that  $U = (U \cap V_{2n}) \oplus (U \cap V_{2l-2n})$ . For such  $U$  the intersection  $U \cap V_{2n}$  is a maximal isotropic subspace of  $V_{2n}$ , and the same holds for  $U \cap V_{2l-2n}$  in  $V_{2l-2n}$ . Therefore every closed

$L_{n,D}$ -orbit  $X := L_{n,D} \cdot U \subset Y$  is isomorphic to the spinor Grassmannian  $GO(n, V_{2n})$ , and its  $Spin(V_{2n})$ -equivariant embedding

$$GO(n, V_{2n}) \xrightarrow{\varphi} GO(l, V_{2l}), \quad \varphi(g \cdot (U \cap V_{2n})) := f_{n,2l}^O(g) \cdot U \quad \text{for } g \in Spin(V_{2n})$$

is a standard extension. Thus statement (ii.1) holds for the closed  $L_{n,D}$ -orbits in  $Y$  and hence (ii) holds for spinor Grassmannians. Proposition 4.2 is proved.  $\square$

*Proof of Theorem 4.1.* The statement is deduced in a straightforward manner from part (i) of Proposition 4.2, by the obvious remark that the tautological embeddings  $GO(m, V_n) \hookrightarrow G(m, V_n)$  and  $GS(m, V_n) \hookrightarrow G(m, V_n)$  are equivariant for the tautological inclusions of groups

$$SO(V_n) \hookrightarrow SL(V_n) \quad \text{and} \quad Sp(V_n) \hookrightarrow SL(V_n),$$

respectively, and any standard extension  $GO(m, V_n) \hookrightarrow GO(m, V_{n+1})$  is equivariant for a homomorphism  $Spin_n \rightarrow Spin_{n+1}$ .  $\square$

Next we consider linear embeddings which factor through a projective space or a standard quadric.

**Proposition 4.3.** *Let  $X \xrightarrow{\varphi} Y$  be a linear embedding of Grassmannians, where  $X$  is not isomorphic to  $GO(m-2, V_{2m})$  for  $m \geq 5$  or to  $GO(m-1, V_{2m+1})$  for  $m \geq 3$ . Assume furthermore that  $\varphi$  factors through a projective space or a quadric. Then  $\varphi$  is equivariant if and only if it is not one of the following embeddings:*

- (a)  $\varphi : X \xrightarrow{\kappa_p} Q \xrightarrow{\sigma} GO(l, V_s) = Y$  for a standard extension  $\sigma$  and some  $p \in I_2(X)$  which is not invariant under any subgroup  $G \subset G_X$  acting transitively on  $X$ .
- (b)  $\varphi : X \xrightarrow{\pi_X} \mathbb{P}(V_X) \xrightarrow{\lambda} GS(1, V_{s-2l}) \xrightarrow{\sigma} GS(l, V_s) = Y$ , where  $2 \leq l < s/2$ ,  $\sigma$  is a standard extension, and  $\lambda$  is a linear embedding such that the restriction of the symplectic form of  $V_{s-2l}$  to  $V_X$  is not  $G$ -invariant for any subgroup  $G \subset G_X$  acting transitively on  $X$ .

*Proof.* Recall that any minimal projective embedding of a Grassmannian  $X$  is  $G_X$ -equivariant. Also, every linear embedding  $\psi : \mathbb{P}(V) \hookrightarrow Y$  of a projective space into a Grassmannian  $Y$  is equivariant, and it is  $SL(V)$ -equivariant if its image is not a standard symplectic projective space on a symplectic Grassmannian. Hence, if  $\varphi$  factors through a projective space which is not a standard symplectic projective space then  $\varphi$  is  $G_X$ -equivariant. If  $\varphi$  factors through a standard symplectic projective space  $GS(1, V_{s-2l})$  on  $Y$  then the condition for equivariance of  $\varphi$  is the invariance of respective symplectic form on  $V_{s-2l}$ , as stated in (b).

Assume that  $\varphi$  factors through a quadric  $Q \subset Y$  but not through a projective space. If  $Y$  is not a spinor Grassmannian, by Corollary 2.9  $Y$  is an orthogonal Grassmannian and  $Q$  is a standard quadric on  $Y$ . By Proposition 2.1 the embedding of  $X$  into  $Q$  is of the form  $\sigma \circ \kappa_p$  for some  $p \in I_2(X)$ . The condition for equivariance of  $\varphi$  is the invariance of  $p$ , as stated in (a). If  $Y$  is a spinor Grassmannian then  $Q$  has dimension at most 6 by Lemma 3.5, and the only linearly embedded Grassmannians in  $Q^6$  are equivariantly embedded quadrics.  $\square$

Let us comment on the occurrence of non-equivariant embeddings. First, recall that in most cases the group  $G_X$  does not admit any proper subgroups acting transitively on  $X$ , and the conditions given in (a) and (b) concern only the group  $G_X$ . The exceptions are only the spinor Grassmannians, and the odd-dimensional projective spaces. In these cases there are indeed linear embeddings which are equivariant under a specific subgroup  $G \subset G_X$ , but not under  $G_X$ . For instance, the minimal projective embedding of  $X = GO(4k+3, V_{8k+6})$  factors through non-degenerate quadrics. None of these quadrics is invariant under  $G_X = Spin(V_{8k+6})$ , but every subgroup  $G \subset G_X$  isomorphic to  $Spin_{8k+5}$  admits such an invariant quadric.

Note that, if  $\varphi : X \hookrightarrow Y$  is a non-equivariant embedding of type (a) or (b) from Proposition 4.3, then the composition of  $\varphi$  with any embedding  $\psi : Y \hookrightarrow Z$  remains non-equivariant if and only if  $\psi$  is a standard extension. All other compositions of  $\varphi$  are equivariant.

On the other hand, if  $\varphi : X \hookrightarrow Y$  and  $\psi : Y \hookrightarrow Z$  are equivariant embeddings then the composition  $\psi \circ \varphi$  is equivariant, unless  $Y$  is a projective space,  $Z \cong GS(l, V_s)$  is a symplectic Grassmannian,  $\psi(Y)$  is a standard symplectic projective space on  $Y$ , and  $\varphi \circ \psi$  is an embedding of type (b) from Proposition 4.3.

In the next proposition we address the equivariance properties of the special embeddings of Grassmannians excluded by the hypothesis of Theorem 4.1.

**Proposition 4.4.** *The embeddings  $\delta_n : GO(n-2, V_{2n}) \hookrightarrow G(2, V_{2n-1})$  and  $\delta_n^1 : GO(n-2, V_{2n-1}) \hookrightarrow G(2, V_{2n-1})$  defined in (8) and (11) are respectively  $Spin(V_{2n})$ - and  $Spin(V_{2n-1})$ -equivariant. Furthermore, the following hold:*

- (i)  $n$  is odd if and only if  $\delta_n$  does not admit factorizations as a composition of two proper equivariant embeddings of Grassmannians;
- (ii)  $n \equiv 2 \pmod{4}$  if and only if  $\delta_n$  factors  $Spin(V_{2n})$ -equivariantly as

$$\delta_n : GO(n-2, V_{2n}) \xrightarrow{\delta_n^S} GS(2, V_{2n-1}) \xrightarrow{\tau} G(2, V_{2n-1});$$

- (iii)  $n \equiv 0 \pmod{4}$  if and only if  $\delta_n$  factors  $Spin(V_{2n+1})$ -equivariantly as

$$\delta_n : GO(n-2, V_{2n}) \xrightarrow{\delta_n^O} GO(2, V_{2n-1}) \xrightarrow{\tau} G(2, V_{2n-1});$$

- (iv)  $n \equiv 1, 2 \pmod{4}$  if and only if  $\delta_n^1$  factors  $Spin(V_{2n-1})$ -equivariantly as

$$\delta_n^1 : GO(n-2, V_{2n-1}) \xrightarrow{\delta_n^{1,S}} GS(2, V_{2n-1}) \xrightarrow{\tau} G(2, V_{2n-1});$$

- (v)  $n \equiv 0, 3 \pmod{4}$  if and only if  $\delta_n^1$  factors  $Spin(V_{2n-1})$ -equivariantly as

$$\delta_n^1 : GO(n-2, V_{2n-1}) \xrightarrow{\delta_n^{1,O}} GO(2, V_{2n-1}) \xrightarrow{\tau} G(2, V_{2n-1}).$$

*Proof.* The existence of an equivariant factorization of  $\delta_n$  through  $GO(2, V_{2n-1})$  or  $GS(2, V_{2n-1})$  is equivalent to the existence of an invariant symmetric or, respectively, symplectic non-degenerate bilinear form on the spin-representation of  $Spin_{2n}$ . The conditions for existence of such invariant bilinear forms and the respective self-duality of spin-representations are well-known, see e.g. [2], [3].  $\square$

**Corollary 4.5.** *Let  $X$  be a Grassmannian.*

- (i) Every linear embedding  $\varphi : X \hookrightarrow G(l, V_s)$  is  $G_X$ -equivariant.
- (ii) Every linear embedding  $\varphi : X \hookrightarrow GO(l, V_{2l})$  is  $G_X$ -equivariant, except possibly if  $X$  is a Lagrangian Grassmannian  $GS(m, V_{2m})$  for  $m \geq 2$ , or  $GO(m-2, V_{2m-\varepsilon})$  for  $5 \leq m$ ,  $\varepsilon \in \{0, 1\}$ .
- (iii) Every linear embedding  $\varphi : X \hookrightarrow GO(l, V_s)$  with  $l < \lfloor \frac{s-1}{2} \rfloor$  is  $G_X$ -equivariant, unless it factors through a non-equivariant embedding into a standard quadric, or through a non-equivariant embedding of the form  $\delta_n^O$  for  $X = GO(n-2, V_{2n})$ , or through a non-equivariant embedding of the form  $\delta_n^{1,O}$  for  $X = GO(n-2, V_{2n-1})$ .
- (iv) Every linear embedding  $\varphi : X \hookrightarrow GS(l, V_s)$  is  $G_X$ -equivariant, unless it factors through a non-equivariant embedding into a standard symplectic projective space, or through a non-equivariant embedding of the form  $\delta_n^S$  for  $X = GO(n-2, V_{2n})$ , or through a non-equivariant embedding of the form  $\delta_n^{1,S}$  for  $X = GO(n-2, V_{2n-1})$ .

*Proof.* For  $X$  not isomorphic to  $GO(n-2, V_{2n})$  or  $GO(n-1, V_{2n+1})$ , the statement follows immediately from Theorem 4.1 and Proposition 4.3. In the two remaining cases the result follows from Theorems 3.3 and 3.4, together with Proposition 4.4.  $\square$

## 5. LINEAR EMBEDDINGS OF LINEAR IND-GRASSMANNIANS

Linear ind-Grassmannians are defined and classified in [9]. In this final section we show that the classification of linear embeddings of Grassmannians  $X \xrightarrow{\varphi} Y$  carries over to linear ind-Grassmannians, with exceptions similar to those in the finite-dimensional case.

**5.1. Linear ind-Grassmannians.** We begin by recalling some basic notions and the classification of ind-Grassmannians.

**Definition 5.1.** *A linear ind-Grassmannian is an ind-variety  $X$  which can be obtained as a direct limit  $X = \lim_{\rightarrow} X_k$  of a sequence of linear embeddings  $\psi_k : X_k \rightarrow X_{k+1}$  of Grassmannians. A sequence  $(X_k, \psi_k)$  with the additional property that all  $\psi_k$  are standard extensions is a **standard exhaustion** of  $\mathbf{X}$ .*

The Picard group of a linear ind-Grassmannian is isomorphic to  $\mathbb{Z}$ , generated by the class of the line bundle  $\mathcal{O}_{\mathbf{X}}(1) := \lim_{\leftarrow} \mathcal{O}_{X_k}(1)$ . Note also that every exhaustion of a linear ind-Grassmannian by Grassmannians is a chain of linear embeddings up to finitely many terms. Indeed, if there were infinitely many non-linear embeddings in an exhaustion, the Picard group of the ind-Grassmannian would be trivial and hence not isomorphic to  $\mathbb{Z}$ .

Let us recall the definitions of the following linear ind-Grassmannians from [9]. Fix a chain of proper inclusions of vector spaces  $\nu_k : V_{n_k} \subset V_{n_{k+1}}$ ,  $k \in \mathbb{Z}_{\geq 1}$ , and denote by  $\{m_k\}$  a non-decreasing sequence of positive integers such that  $\{n_k - m_k\}$  is also a non-decreasing sequence of positive integers.

(A)  $\mathbf{G}(m) := \lim_{\rightarrow} G(m, V_{n_k})$  is defined as the direct limit of the chain

$$\dots \hookrightarrow G(m, V_{n_k}) \xrightarrow{\sigma_k} G(m, V_{n_{k+1}}) \hookrightarrow \dots$$

of strict standard extensions associated to the inclusions  $\nu_k$ , where we assume  $m := m_k < n_1$  for all  $k$ .

(B)  $\mathbf{G}(\infty) := \lim_{\rightarrow} G(m_k, V_{n_k})$ , the **Sato Grassmannian** [10], is defined as the direct limit of an arbitrary chain of standard extensions

$$\dots \hookrightarrow G(m_k, V_{n_k}) \xrightarrow{\sigma_k} G(m_{k+1}, V_{n_{k+1}}) \hookrightarrow \dots$$

associated to the inclusions  $\nu_k$ , under the assumption that  $\lim_{k \rightarrow \infty} m_k = \infty = \lim_{k \rightarrow \infty} (n_k - m_k)$ .

(C)  $\mathbf{GO}(m, \infty) := \lim_{\rightarrow} GO(m, V_{n_k})$  is defined as the direct limit of the chain

$$\dots \hookrightarrow GO(m, V_{n_k}) \xrightarrow{\sigma_k} GO(m, V_{n_{k+1}}) \hookrightarrow \dots$$

of standard extensions associated to the inclusions  $\nu_k$ , which here are assumed to be non-degenerate inclusions of orthogonal vector spaces, and where  $m := m_k < \lfloor \frac{n_1}{2} \rfloor$  for all  $k$ .

(D)  $\mathbf{GO}(\infty, \infty) := \lim_{\rightarrow} GO(m_k, V_{n_k})$  is defined as the direct limit of an arbitrary chain of standard extensions

$$\dots \hookrightarrow GO(m_k, V_{n_k}) \xrightarrow{\sigma_k} GO(m_{k+1}, V_{n_{k+1}}) \hookrightarrow \dots$$

compatible with the inclusions  $\nu_k$ , which are assumed to be non-degenerate inclusions of orthogonal vector spaces, and where  $\lfloor \frac{n_k}{2} \rfloor - m_k$  is a non-decreasing sequence of positive integers,  $\lim_{k \rightarrow \infty} m_k = \infty$  and  $\lim_{k \rightarrow \infty} (\lfloor \frac{n_k}{2} \rfloor - m_k) = \infty$ .

(E)  $\mathbf{GO}^0(\infty, m) := \lim_{\rightarrow} GO(m_k, V_{n_k})$ , for  $m \geq 2$ , is defined as the direct limit of an arbitrary chain of standard extensions

$$\dots \hookrightarrow GO(m_k, V_{n_k}) \xrightarrow{\sigma_k} GO(m_{k+1}, V_{n_{k+1}}) \hookrightarrow \dots$$

compatible with the inclusions  $\nu_k$ , which are assumed to be non-degenerate inclusions of orthogonal vector spaces, and where  $n_k$  is even for all  $k$ ,  $\frac{n_k}{2} - m_k$  is a non-decreasing sequence of positive integers,  $\lim_{k \rightarrow \infty} m_k = \infty$  and  $\lim_{k \rightarrow \infty} (\frac{n_k}{2} - m_k) = m$ .

- (F)  $\mathbf{GO}^1(\infty, m) := \lim_{\rightarrow} GO(m_k, V_{n_k})$  for  $m \geq 0$  is defined as the direct limit of an arbitrary chain of standard extensions

$$\dots \hookrightarrow GO(m_k, V_{n_k}) \xrightarrow{\sigma_k} GO(m_{k+1}, V_{n_{k+1}}) \hookrightarrow \dots$$

compatible with the inclusions  $\nu_k$ , which are assumed to be non-degenerate inclusions of orthogonal vector spaces, and where  $n_k$  is odd for all  $k$ ,  $\lfloor \frac{n_k}{2} \rfloor - m_k$  is a non-decreasing sequence of positive integers,  $\lim_{k \rightarrow \infty} m_k = \infty$  and  $\lim_{k \rightarrow \infty} (\lfloor \frac{n_k}{2} \rfloor - m_k) = m$ .

- (G)  $\mathbf{GS}(m, \infty) := \lim_{\rightarrow} GS(m, V_{n_k})$  is defined as the direct limit of the chain

$$\dots \hookrightarrow GS(m, V_{n_k}) \xrightarrow{\sigma_k} GS(m, V_{n_{k+1}}) \hookrightarrow \dots$$

of standard extensions associated to the inclusions  $\nu_k$ , which are now assumed to be non-degenerate inclusions of symplectic vector spaces, and where  $m := m_k < \lfloor \frac{n_1}{2} \rfloor$  for all  $k$ .

- (H)  $\mathbf{GS}(\infty, \infty) := \lim_{\rightarrow} GS(m_k, V_{n_k})$  is defined as an analogue of case (D) for symplectic Grassmannians.

- (I)  $\mathbf{GS}(\infty, m) := \lim_{\rightarrow} GS(m_k, V_{n_k})$  for  $m \geq 0$  is defined as the direct limit of an arbitrary chain of standard extensions

$$\dots \hookrightarrow GS(m_k, V_{n_k}) \xrightarrow{\sigma_k} GS(m_{k+1}, V_{n_{k+1}}) \hookrightarrow \dots$$

compatible with the inclusions  $\nu_k$ , which are assumed to be non-degenerate inclusions of symplectic vector spaces, and where under the assumption that  $n_k$  is even for all  $k$ ,  $\frac{n_k}{2} - m_k$  is a non-decreasing sequence of positive integers,  $\lim_{k \rightarrow \infty} m_k = \infty$  and  $\lim_{k \rightarrow \infty} (\frac{n_k}{2} - m_k) = m$ .

**Proposition 5.1.** ([9, Lemma 4.3]) *Each of the ind-Grassmannians defined above depends up to isomorphism only on the respective limits  $\lim_{k \rightarrow \infty} m_k$ ,  $\lim_{k \rightarrow \infty} (n_k - m_k)$  and  $\lim_{k \rightarrow \infty} (\lfloor \frac{n_k}{2} \rfloor - m_k)$ .*

Note that all ind-Grassmannians defined above are defined only up to isomorphism. Furthermore, the ind-Grassmannians  $\mathbf{G}(m)$  admit an alternative construction as  $\mathbf{G}(m) \cong \lim_{\rightarrow} G(m_k, V_{n_k})$  where  $n_k - m_k = m$  for all  $k$ .

The ind-Grassmannian  $\mathbf{G}(1)$  is the **projective ind-space**  $\mathbb{P}^\infty$ , and is isomorphic to  $\mathbf{GS}(1, \infty)$ . The ind-Grassmannian  $\mathbf{GO}(1, \infty)$  is the **ind-quadric**  $Q^\infty$ .

**Theorem 5.2.** ([9, Theorem 2]) *Every linear ind-Grassmannian is isomorphic as an ind-variety to one of the ind-Grassmannians  $\mathbf{G}(m)$  for  $m \geq 1$ ,  $\mathbf{G}(\infty)$ ,  $\mathbf{GO}(m, \infty)$  for  $m \geq 1$ ,  $\mathbf{GO}^0(\infty, m)$  for  $m \geq 2$ ,  $\mathbf{GO}^1(\infty, m)$  for  $m \geq 0$ ,  $\mathbf{GO}(\infty, \infty)$ ,  $\mathbf{GS}(m, \infty)$  for  $m \geq 2$ ,  $\mathbf{GS}(\infty, m)$  for  $m \geq 0$ ,  $\mathbf{GS}(\infty, \infty)$ , and the latter are pairwise non-isomorphic.*

We should point out that, despite Theorem 5.2, some linear ind-Grassmannians admit exhaustions which are not standard. For instance, the Sato Grassmannian  $\mathbf{G}(\infty)$  can be obtained as a direct limit of a chain

$$(18) \quad \dots \hookrightarrow GS(m_{2l}, V_{n_{2l}}) \xrightarrow{\varphi_{2l}} GO(m_{2l+1}, V_{n_{2l+1}}) \xrightarrow{\varphi_{2l+1}} GS(m_{2l+2}, V_{n_{2l+2}}) \hookrightarrow \dots$$

where each  $\varphi_k$  is a mixed combination of standard and isotropic extensions. Indeed, a chain of embeddings of the form (18) is easy to define under the assumption that  $\lim_{s \rightarrow \infty} m_s = \infty = \lim_{s \rightarrow \infty} (\frac{n_s}{2} - m_s)$ . Then the direct limit of such a chain will be isomorphic to the direct limit of the

subchain consisting of ordinary Grassmannians only, and the latter direct limit is necessarily isomorphic to  $\mathbf{G}(\infty)$  since the embeddings of this chain do not factor through projective spaces.

We now give an alternative definition of the linear ind-Grassmannians. One can think of the definitions (A)-(I) as local, while the definition below is global and yields a well-defined ind-variety rather than an isomorphism class of ind-varieties.

Let  $V$  be a fixed vector space of infinite countable dimension. Let  $W \subset V$  be a subspace and  $E \subset V$  be a basis of  $V$  such that  $E \cap W$  is a basis of  $W$ . Let  $\mathbf{G}(W, E, V)$  denote the set of subspaces  $U \subset V$  which are  **$E$ -commensurable** with  $W$ , i.e., satisfy:

- there exists a finite-dimensional subspace  $T_U \subset V$  such that  $U \subset W + T_U$ ,  $W \subset U + T_U$  and  $\dim(U \cap T_U) = \dim(W \cap T_U)$ ;
- $\text{codim}_U \text{span}(E \cap U) < \infty$ .

Let now  $V$  be endowed with a symmetric or skew-symmetric non-degenerate bilinear form  $\omega$ . A basis  $E$  of  $V$  is **isotropic** if it admits an involution  $i_E : E \rightarrow E$  with at most one fixed point, and such that  $\omega(e, e') = 0$  for  $e, e' \in E$  unless  $e' = i_E(e)$ . For a fixed isotropic subspace  $W \subset V$  and an isotropic basis  $E \subset V$  containing a basis of  $W$ , we denote by  $\mathbf{GO}(W, E, V)$ , or respectively  $\mathbf{GS}(W, E, V)$ , the set of isotropic subspaces of  $V$  which are  $E$ -commensurable with  $W$ .

**Theorem 5.3.** ([1]) *The sets  $\mathbf{G}(W, E, V)$ ,  $\mathbf{GO}(W, E, V)$ ,  $\mathbf{GS}(W, E, V)$  admit structures of projective ind-varieties isomorphic to linear ind-Grassmannians as follows:*

$$\begin{aligned} \mathbf{G}(W, E, V) &\cong \mathbf{G}(\min\{\dim W, \text{codim}_V W\}) , \\ \mathbf{GO}(W, E, V) &\cong \begin{cases} \mathbf{GO}(\dim W, \infty) & \text{for } \text{codim}_U W = \infty , \\ \mathbf{GO}^\varepsilon(\infty, \text{codim}_U W) & \text{for } 0 < \text{codim}_U W < \infty , \\ \mathbf{GO}^1(\infty, \text{codim}_U W) & \text{for } \text{codim}_U W = 0 , \end{cases} \\ \mathbf{GS}(W, E, V) &\cong \mathbf{GS}(\dim W, \text{codim}_U W) , \end{aligned}$$

where  $U \subset V$  is a maximal isotropic subspace containing  $W$  and spanned by  $E \cap U$ , and  $\varepsilon \in \{0, 1\}$  is the number of fixed points of the involution  $\iota_E$ .

Let  $\mathbf{X}$  be a linear ind-Grassmannian with a fixed standard exhaustion  $\mathbf{X} = \lim_{\rightarrow} X_k$ . Then a triple  $(W, E, V)$  yielding an ind-Grassmannian  $\mathbf{G}(W, E, V)$  isomorphic to  $\mathbf{X}$  is obtained by setting  $V := \lim_{\rightarrow} V_{n_k}$  and  $W := \lim_{\rightarrow} U_k$ , where  $U_1 \in X_1$  is fixed arbitrarily,  $U_{k+1} := \psi_k(U_k) \in X_{k+1}$  for  $k \geq 1$ , and  $E \subset V$  is a basis such that  $E \cap V_{n_k}$  is a basis of  $V_{n_k}$  and  $E \cap U_k$  is a basis of  $U_k$ . Moreover, as a set and ind-variety,  $\mathbf{G}(W, E, V)$  depends not on the entire basis  $E$  but only on the intersection  $W \cap E$  up to changing finitely many basis vectors in  $W \cap E$ . In particular, if  $\dim W = m < \infty$  then  $\mathbf{G}(W, E, V)$  does not depend on  $E$  and depends only on  $m$  and not on  $W$ . In this case we may write  $\mathbf{G}(m, V)$ , and  $\mathbb{P}(V)$  for  $m = 1$ .

Let us remark that **the spinor ind-Grassmannian**  $\mathbf{GO}^1(\infty, 0)$  admits two alternative realizations as  $\mathbf{GO}(W, E, W \oplus W_* \oplus V_1) \cong \lim_{\rightarrow} \mathbf{GO}(k, V_{2k+1})$  and  $\mathbf{GO}(W, E, W \oplus W_*) \cong \lim_{\rightarrow} \mathbf{GO}(k, V_{2k})$ , where  $W_* = \iota_E(E \cap W)$  and  $V_1 = W^\perp \cap (W_*)^\perp$ . In some constructions it will be convenient to use the notation  $\mathbf{GO}^0(\infty, 0)$  for the latter realization, noting that  $\mathbf{GO}^1(\infty, 0) \cong \mathbf{GO}^0(\infty, 0)$ .

## 5.2. Linear embeddings.

**Definition 5.2.** *Let  $\mathbf{X}$  and  $\mathbf{Y}$  be linear ind-Grassmannians. A morphism of ind-varieties  $\mathbf{X} \xrightarrow{\varphi} \mathbf{Y}$  is a **linear embedding** if there exist standard exhaustions  $\mathbf{X} = \lim_{\rightarrow} X_k$  and  $\mathbf{Y} = \lim_{\rightarrow} Y_k$  such that  $\varphi$  is equal to the direct limit  $\lim_{\rightarrow} \varphi_k$  of a sequence of linear embeddings  $X_k \xrightarrow{\varphi_k} Y_k$ .*

If  $\mathbf{X} \xrightarrow{\varphi} \mathbf{Y}$  is a linear embedding as above then  $\mathcal{O}_{\mathbf{X}}(1) \cong \varphi^* \mathcal{O}_{\mathbf{Y}}(1)$  holds.

Note that it is possible to obtain an isomorphism of linear ind-Grassmannians as a direct limit of proper linear embeddings  $\varphi_k$  of Grassmannians. Indeed, consider some linear ind-Grassmannian  $\mathbf{X} = \varinjlim X_k$  with defining chain of embeddings  $\psi_k : X_k \hookrightarrow X_{k+1}$ . By setting  $Y_k := X_{k+1}$  and  $\varphi_k := \psi_k$ , we obtain a sequence of linear embeddings  $X_k \xrightarrow{\varphi_k} Y_k$  which induces the identity map on  $\mathbf{X} = \varinjlim X_k = \varinjlim Y_k$ . On the other hand, there are proper linear embeddings between isomorphic linear ind-Grassmannians. In particular, any proper inclusion of countable-dimensional vector spaces  $V' \subset V$  induces a linear embedding  $\mathbb{P}(V') \hookrightarrow \mathbb{P}(V)$ .

Next we define several types of embeddings of linear ind-Grassmannians, generalizing the constructions from the finite-dimensional case. We use the realizations provided by Theorem 5.3.

Note first that every linear ind-Grassmannian  $\mathbf{X}$  admits a linear embedding into a projective ind-space. Indeed, there are natural inclusions  $V_{X_k} \subset V_{X_{k+1}}$  for a given standard exhaustion  $\mathbf{X} = \varinjlim X_k$ , and we can set  $V_{\mathbf{X}} := \varinjlim V_{X_k}$ . Then a linear projective embedding

$$\pi_{\mathbf{X}} : \mathbf{X} \hookrightarrow \mathbb{P}(V_{\mathbf{X}})$$

is obtained as the direct limit  $\pi_{\mathbf{X}} := \varinjlim \pi_{X_k}$  of the minimal projective embeddings  $\pi_{X_k} : X_k \hookrightarrow \mathbb{P}(V_{X_k})$ .

An embedding  $\mathbf{G}(m) \xrightarrow{\varphi} \mathbf{G}(l)$ , where  $m, l \in \mathbb{N} \cup \{\infty\}$ , is a **standard extension** if  $\varphi$  can be expressed as

$$(19) \quad \begin{array}{ccc} \varphi : \mathbf{G}(m) \cong \mathbf{G}(W, E, V) & \hookrightarrow & \mathbf{G}(\tilde{W}, \tilde{E}, \tilde{V}) \cong \mathbf{G}(l) \\ U & \mapsto & U \oplus W' \end{array}$$

for some isomorphisms  $\mathbf{G}(m) \cong \mathbf{G}(W, E, V)$  and  $\mathbf{G}(l) \cong \mathbf{G}(\tilde{W}, \tilde{E}, \tilde{V})$ , together with an inclusion of infinite-dimensional vector spaces  $V \subset \tilde{V}$  such that  $E = \tilde{E} \cap V$ , a decomposition  $\tilde{V} = V \oplus V'$ , and a subspace  $W' \subset V'$  with basis  $\tilde{E} \cap W'$  such that  $\tilde{W} = W \oplus W'$ . Note that  $m \leq l$  necessarily holds in the above situation.

A standard extension  $\mathbf{GS}(m, c) \xrightarrow{\varphi} \mathbf{GS}(l, d)$  is defined by the same formula  $U \mapsto U \oplus W'$  as above, under the additional requirements that the inclusion  $V \subset \tilde{V}$  is non-degenerate, the subspaces  $W$  and  $\tilde{W}$  are isotropic, the bases  $E, \tilde{E}$  are isotropic. The same applies for standard extensions  $\mathbf{GO}(m, \infty) \xrightarrow{\varphi} \mathbf{GO}(l, \infty)$ , as well as for  $\mathbf{GO}^\varepsilon(\infty, m) \xrightarrow{\varphi} \mathbf{GO}^{\varepsilon'}(\infty, l)$  with  $\varepsilon, \varepsilon' \in \{0, 1\}$ . In the latter case we require  $l = 0$  whenever  $m = 0$ .

It is straightforward to verify that every standard extension of linear ind-Grassmannians is a linear embedding. Furthermore, a standard extension between non-spinor ind-Grassmannians is a proper embedding if and only if the inclusion  $V \subset \tilde{V}$  is proper.

A **tautological embedding** of  $\mathbf{GS}(m, \infty)$  (and analogously of  $\mathbf{GO}(m, \infty)$ ) into  $\mathbf{G}(m)$  is defined for any isomorphism  $\mathbf{GS}(m, \infty) \cong \mathbf{GS}(W, E, V)$  (where  $\dim W = m$ ) as

$$\tau_{\mathbf{GS}(W, E, V)} : \mathbf{GS}(m, \infty) \cong \mathbf{GS}(W, E, V) \hookrightarrow \mathbf{G}(W, E, V) \cong \mathbf{G}(m) . \\ U \mapsto U$$

This embedding is linear for all  $m > 0$ .

A **tautological embedding** of  $\mathbf{GO}^\varepsilon(\infty, m)$  into  $\mathbf{G}(\infty)$  is defined for any isomorphism  $\mathbf{GO}^\varepsilon(\infty, m) \cong \mathbf{GO}(W, E, V)$  (where  $\dim W = \infty$  and  $W$  has codimension  $m$  in a maximal isotropic subspace of  $V$ ) as

$$\tau_{\mathbf{GO}(W, E, V)} : \mathbf{GO}^\varepsilon(\infty, m) \cong \mathbf{GO}(W, E, V) \hookrightarrow \mathbf{G}(W, E, V) \cong \mathbf{G}(\infty) . \\ U \mapsto U$$

This embedding is linear for  $m > 0$ .

A **tautological embedding** of  $\mathbf{GS}(\infty, m)$  into  $\mathbf{G}(\infty)$  is defined analogously, and is linear for all  $m \geq 0$ .

An **isotropic extension** of  $\mathbf{G}(m)$  to  $\mathbf{GS}(m, \infty)$ , and to  $\mathbf{GS}(\infty, n)$  for  $n \geq m$ , is defined for a given isotropic subspace  $V \subset \tilde{V}$  and an isotropic basis  $F \subset \tilde{V}$  such that  $E := V \cap F$  is a basis of  $V$ , as an embedding

$$\begin{array}{ccc} \iota : \mathbf{G}(m) \cong \mathbf{G}(W, E, V) & \hookrightarrow & \mathbf{GS}(W, F, \tilde{V}) \cong \mathbf{GS}(m, \infty) \\ U & \mapsto & U \end{array}$$

where  $W \subset V$  is a subspace of dimension  $m$  and infinite codimension, and respectively as

$$\begin{array}{ccc} \iota : \mathbf{G}(m) \cong \mathbf{G}(V_\infty, E, V) & \hookrightarrow & \mathbf{GS}(V_\infty, F, \tilde{V}) \cong \mathbf{GS}(\infty, n) \\ U & \mapsto & U \end{array}$$

where  $V_\infty \subset V$  is a subspace of codimension  $m$  in  $V$  and codimension  $n$  in a maximal isotropic subspace of  $\tilde{V}$ .

Isotropic extensions of  $\mathbf{G}(m)$  to  $\mathbf{GO}(m, \infty)$ , or to  $\mathbf{GO}^\varepsilon(\infty, n)$  with  $n \geq m$ , are defined analogously. All isotropic extensions are linear. Note that the ind-Grassmannians  $\mathbf{GS}(\infty, 0)$  and  $\mathbf{GO}^\varepsilon(\infty, 0)$  are not targets of isotropic extensions.

Composing tautological embeddings, standard extensions and isotropic extensions, we define **(mixed) combinations of standard and isotropic extensions** by analogy with the finite dimensional case.

A **projective ind-space on** (or a **projective space on**) an ind-Grassmannian  $\mathbf{X}$  is the image of a linear embedding  $\mathbb{P}^\infty \hookrightarrow \mathbf{X}$  (respectively,  $\mathbb{P}^k \hookrightarrow \mathbf{X}$  for some  $k$ ). An **ind-quadric on  $\mathbf{X}$**  is the image of a linear embedding  $Q^\infty \hookrightarrow \mathbf{X}$ . For  $\mathbf{X} \cong \mathbf{GO}(m, \infty)$  with  $m \in \mathbb{N} \cup \{\infty\}$ , a **standard ind-quadric on  $\mathbf{X}$**  is the image of a standard extension  $Q^\infty \hookrightarrow \mathbf{X}$ . For  $\mathbf{X} \cong \mathbf{GS}(m, \infty)$  with  $m \in \mathbb{N} \cup \{\infty\}$ , a **standard symplectic projective ind-space on  $\mathbf{X}$**  is the image of a standard extension  $\mathbb{P}^\infty \cong \mathbf{GS}(1, \infty) \hookrightarrow \mathbf{X}$ .

**Proposition 5.4.** *Let  $\mathbf{X}$  and  $\mathbf{Y}$  be linear ind-Grassmannians.*

- (i) *There exists a linear embedding  $\mathbf{X} \xrightarrow{\varphi} \mathbf{Y}$  which factors through a projective ind-space, if and only if  $\mathbf{Y}$  is not isomorphic to  $\mathbf{GS}(\infty, 0)$ .*
- (ii) *Suppose  $\mathbf{X}$  is not a projective ind-space or an ind-quadric, and fix a standard exhaustion  $\mathbf{X} = \lim_{\rightarrow} X_k$  such that none of the Grassmannians  $X_k$  is a projective space or a quadric.*

*Let  $\mathbf{X} \xrightarrow{\varphi} \mathbf{Y}$  be a linear embedding obtained as the direct limit of a sequence  $X_k \xrightarrow{\varphi_k} Y_k$  for a suitable standard exhaustion  $\mathbf{Y} = \lim_{\rightarrow} Y_k$ . Then  $\varphi$  factors through a projective ind-space or through a standard ind-quadric if and only if the embedding  $\varphi_1$  factors through a projective space or a standard ind-quadric.*

*Proof.* Part (i) follows from the fact that every linear ind-Grassmannian admits a linear embedding into a projective ind-space, together with the observation that there is an infinite-dimensional projective ind-space on every linear ind-Grassmannian except  $\mathbf{GS}(\infty, 0)$ . The maximal projective spaces on  $\mathbf{GS}(\infty, 0)$  are projective lines.

For part (ii), suppose first that  $\mathbf{X}$  is not isomorphic to  $\mathbf{GS}(\infty, 0)$ . We may also assume that  $\mathbf{Y}$  is not a projective ind-space, an ind-quadric, or  $\mathbf{GS}(0, \infty)$ . Let us recall that on a Grassmannian  $Z$  which is not a projective space, a quadric, or  $GS(l, V_{2l})$ , there are two irreducible families  $\mathcal{F}_1(Z), \mathcal{F}_2(Z)$  of maximal projective spaces, unless  $Z$  is a non-spinor orthogonal Grassmannians which has one family  $\mathcal{F}_1(Z)$  of maximal standard quadrics and one family  $\mathcal{F}_2(Z)$  of maximal projective spaces not contained in a quadric. If  $\psi : Z_1 \rightarrow Z_2$  is a linear embedding of such Grassmannians  $Z_1, Z_2$ , and  $\psi$  does not factor through a projective space or through a standard quadric, then  $\psi$  separates the families  $\mathcal{F}_1$  and  $\mathcal{F}_2$ , i.e., after possible relabelling of the families  $\mathcal{F}_1(Z_2), \mathcal{F}_2(Z_2)$  we can assume that  $\psi$  maps spaces from  $\mathcal{F}_j(Z_1)$  to spaces from  $\mathcal{F}_j(Z_2)$  for  $j = 1, 2$ . Note in addition that a projective space of dimension  $l \geq 3$  on  $Z$  is contained in a unique maximal projective space or in a unique maximal standard quadric on  $Z$ .

If  $\varphi_1$  factors through a projective space  $P \subset Y_1$  then  $3 \leq \dim X_1 \leq \dim P$  holds, and hence  $P$  determines a unique maximal projective space or a unique maximal standard quadric  $M_k \subset Y_k$  for each  $k \geq 1$ . Furthermore, either each  $M_k$  is a projective space, or each  $M_k$  is a quadric. We clearly have  $M_k \subset M_{k+1}$ , and thus  $P$  determines a unique maximal projective space or maximal standard quadric  $M \subset \mathbf{Y}$ . We claim that  $\varphi_k(X_k) \subset M_k$ . Indeed, for every  $k \geq 1$  there exists  $F_j^k \in \mathcal{F}_j(X_k)$  for  $j = 1, 2$  such that  $F_j^k \cap X_1 \in \mathcal{F}_j(X_1)$ . Then necessarily  $\varphi_k(F_j^k) \subset M_k$ . This implies that  $\varphi_k$  sends both families  $\mathcal{F}_1(X_k), \mathcal{F}_2(X_k)$  to one family, say  $\mathcal{F}_1(Y_k)$ , on  $Y_k$ . Consequently, we have  $\varphi(\mathbf{X}) \subset M$ .

Suppose now that  $\varphi_1$  does not factor through a projective space or through a standard quadric. Then any two spaces  $F_j \in \mathcal{F}_j(X_1)$ ,  $j = 1, 2$ , determine unique sequences  $\tilde{F}_j^k \in \mathcal{F}_j(Y_k)$  for  $k \geq 1$  such that  $\tilde{F}_j^k \cap \varphi(\mathbf{X}) \geq 2$  for all  $k$  and for  $j = 1, 2$ . Thus the image  $\varphi(\mathbf{X})$  is not contained in a projective ind-space or an ind-quadric on  $\mathbf{Y}$ .

It remains to consider the case  $X = \mathbf{GS}(\infty, 0)$ . Due to Theorems 2.4 and 2.5, every linear embedding of  $GS(m, V_{2m})$  which does not factor through a projective space is a standard extension  $GS(m, V_{2m}) \hookrightarrow GS(l, V_{2l})$  or factors through the tautological embedding  $GS(m, V_{2m}) \hookrightarrow G(m, V_{2m})$ . The case of a standard extension is excluded since  $\mathbf{Y} \not\cong \mathbf{GS}(\infty, 0)$ . The remaining options allow us to reduce the proof to the case of ordinary ind-Grassmannians.  $\square$

For the following proposition, we recall that every linear embedding of a Grassmannian  $X$  into a quadric factors through one of the embeddings  $\kappa_p$  for  $p \in I_2(X)$  described in Proposition 2.1. Also, every linear embedding  $\psi : X \hookrightarrow Y$  of Grassmannians induces a linear embedding of projective spaces  $\hat{\psi} : \mathbb{P}(V_X) \hookrightarrow \mathbb{P}(V_Y)$  as in (1), and hence a pullback  $\hat{\psi}^* : \mathbb{C}[V_Y] \rightarrow \mathbb{C}[V_X]$  on polynomial functions. Furthermore, we have  $\hat{\psi}^*(I_2(Y)) \subset I_2(X)$ . For a linear ind-Grassmannian  $\mathbf{X}$  with a standard exhaustion  $X_k \xrightarrow{\psi_k} X_{k+1}$ , we obtain the inverse limit  $I_2(\mathbf{X}) = \lim_{\leftarrow} I_2(X_k)$  consisting of sequences  $p_k \in I_2(X_k)$  satisfying  $p_k = \hat{\psi}_k^* p_{k+1}$ .

**Proposition 5.5.** *Let  $\mathbf{X} \xrightarrow{\varphi} Q^\infty$  be a linear embedding. If  $\varphi$  does not factor through a projective ind-space, then  $\varphi$  factors as*

$$\mathbf{X} \xrightarrow{\kappa_{\mathbf{p}}} Q^\infty \xrightarrow{\sigma} Q^\infty$$

where  $\sigma$  is a standard extension,  $\mathbf{p} \in I_2(\mathbf{X})$  is a nonzero element, and  $\kappa_{\mathbf{p}} := \lim_{\rightarrow} \kappa_{p_k}$ .

*Proof.* Let  $\tilde{\psi}_k : Q^{s_k} \hookrightarrow Q^{s_{k+1}}$  be standard extensions defining  $\mathbf{Y} := Q^\infty$ , such that the embedding  $\varphi$  is the direct limit of a sequence of linear embeddings  $X_k \xrightarrow{\varphi_k} Q^{s_k}$ , none of which factors through a projective space. By Proposition 2.1, each  $\varphi_k$  factors as  $\varphi_k = \sigma_k \circ \kappa_{p_k}$  for a standard extension  $\sigma_k : Q^{n_k} \hookrightarrow Q^{s_k}$  and a nonzero  $p_k \in I_2(X_k)$ . The relation  $p_k = \hat{\psi}_k^* p_{k+1}$  follows from the observation that  $\varphi_{k+1} \circ \psi_k = \tilde{\psi}_k \circ \varphi_k$ , since  $p_k$  is equal to the restriction to  $V_{X_k}$  of a non-degenerate quadric form defining  $Q^{s_k}$ . Thus we can form the element  $\mathbf{p} := \lim_{\leftarrow} p_k \in I_2(\mathbf{X})$ .

The quadrics  $Q^{n_k}$  satisfy

$$\tilde{\psi}_k(\varphi_k(X_k)) \subset \tilde{\psi}_k(\sigma_k(Q^{n_k})) \subset \sigma_{k+1}(Q^{n_{k+1}}),$$

and we obtain a chain of standard extensions  $\eta_k : Q^{n_k} \hookrightarrow Q^{n_{k+1}}$  fitting into a commutative diagram

$$\begin{array}{ccccccc} \varphi_{k+1} : & X_{k+1} & \xrightarrow{\kappa_{p_{k+1}}} & Q^{n_{k+1}} & \xrightarrow{\sigma_{k+1}} & Q^{s_{k+1}} & \\ & \psi_k \uparrow & & \eta_k \uparrow & & \tilde{\psi}_k \uparrow & \\ \varphi_k : & X_k & \xrightarrow{\kappa_{p_k}} & Q^{n_k} & \xrightarrow{\sigma_k} & Q^{s_k} & \end{array} .$$

We can now conclude that the embedding  $\varphi$  factors through the ind-quadric  $\mathbf{Z} = \lim_{\rightarrow} Q^{n_k} \cong Q^\infty$  defined by the embeddings  $\eta_k$ , as

$$\varphi : \mathbf{X} \xrightarrow{\kappa_{\mathbf{p}}} \mathbf{Z} \xrightarrow{\sigma} \mathbf{Y}$$

where  $\kappa_{\mathbf{p}} = \lim_{\rightarrow} \kappa_{p_k}$  and  $\sigma = \lim_{\rightarrow} \sigma_k$ . □

The next proposition requires some preparation. Let  $V$  be an orthogonal space,  $W \subset V$  be a maximal isotropic subspace, and  $E \subset V$  be an isotropic basis containing a basis of  $W$ . We assume that the involution  $\iota_E$  has no fixed points, so that  $V = W \oplus W_*$  where  $W_* = \text{span}(\iota_E(E \cap W))$ . Then  $\mathbf{S} := \mathbf{GO}(W, E, V)$  is isomorphic to the spinor ind-Grassmannian  $\mathbf{GO}^0(\infty, 0)$ . Let  $\mathbf{S} \xrightarrow{\pi_{\mathbf{S}}} \mathbb{P}(V_{\mathbf{S}})$  be the minimal projective embedding of  $\mathbf{S}$ .

Let  $W^{(-2)} \subset W$  be a fixed subspace of codimension 2, spanned by  $E \cap W^{(-2)}$ . Then  $\mathbb{P}_{\mathbf{S}, U}^1 := \{U' \in \mathbf{S} : U \subset U'\}$  for  $U \in \mathbf{GO}^0(W^{(-2)}, E, V)$  is a projective line on  $\mathbf{S}$ , and every projective line on  $\mathbf{S}$  has this form for some (unique)  $U$ . Thus the set of projective lines on  $\mathbf{S}$  admits a structure of a linear ind-Grassmannian isomorphic to  $\mathbf{GO}^0(\infty, 2)$ .

**Proposition 5.6.** *The map*

$$\begin{aligned} \delta^0 : \mathbf{GO}^0(\infty, 2) \cong \mathbf{GO}(W^{(-2)}, E, V) &\hookrightarrow \mathbf{G}(2, V_{\mathbf{S}}) \cong \mathbf{G}(2) . \\ U &\mapsto \pi_{\mathbf{S}}(\mathbb{P}_{\mathbf{S}, U}^1) \end{aligned}$$

is a proper linear embedding of linear ind-Grassmannians. The embedding  $\delta^0$  admits two factorizations:

$$\mathbf{GO}^0(\infty, 2) \xrightarrow{\delta_{\mathbf{S}}^0} \mathbf{GS}(2, \infty) \xrightarrow{\tau_{\mathbf{S}}} \mathbf{G}(2) \quad , \quad \mathbf{GO}^0(\infty, 2) \xrightarrow{\delta_{\mathbf{O}}^0} \mathbf{GO}(2, \infty) \xrightarrow{\tau_{\mathbf{O}}} \mathbf{G}(2) \quad ,$$

where  $\tau_{\mathbf{S}}, \tau_{\mathbf{O}}$  are the respective tautological embeddings.

*Proof.* The linearity of the map  $\delta^0$  follows from a local expression as a direct limit  $\delta^0 = \lim_{\rightarrow} \delta_n$ , where  $\delta_n : \mathbf{GO}(n-2, V_{2n}) \hookrightarrow \mathbf{G}(2, V_{2n-1})$  are the embeddings defined in (8).

The embeddings  $\delta_{\mathbf{S}}^0$  and  $\delta_{\mathbf{O}}^0$  are obtained as direct limits  $\delta_{\mathbf{S}}^0 = \lim_{\rightarrow} \delta_{4k+2}^{\mathbf{S}}$  and  $\delta_{\mathbf{O}}^0 = \lim_{\rightarrow} \delta_{4k}^{\mathbf{O}}$  of the embeddings given in Proposition 4.4. □

The compositions of a standard extension  $\mathbf{GO}^1(\infty, 1) \xrightarrow{\sigma} \mathbf{GO}^0(\infty, 2)$  with the embeddings  $\delta^0, \delta_{\mathbf{S}}^0, \delta_{\mathbf{O}}^0$  yield linear embeddings

$$(20) \quad \begin{aligned} \delta^1 &= \delta^0 \circ \sigma : \mathbf{GO}^1(\infty, 1) \hookrightarrow \mathbf{G}(2) \quad , \\ \delta_{\mathbf{S}}^1 &= \delta_{\mathbf{S}}^0 \circ \sigma : \mathbf{GO}^1(\infty, 1) \hookrightarrow \mathbf{GS}(2, \infty) \quad , \\ \delta_{\mathbf{O}}^1 &= \delta_{\mathbf{O}}^0 \circ \sigma : \mathbf{GO}^1(\infty, 1) \hookrightarrow \mathbf{GO}(2, \infty) \quad . \end{aligned}$$

Next we describe the inductive limit of the embeddings  $\theta_m^k$  defined in Subsection 3.1. Let  $V$  be an orthogonal vector space of infinite countable dimension,  $W \subset V$  be a maximal isotropic subspace,  $W^{(-k)} \subset W$  be a subspace of codimension  $k$ ,  $E$  be a basis of  $W$  containing a basis of  $W^{(-k)}$ ,  $\tilde{E}$  be an isotropic basis of  $V$  containing  $E$ , and  $E_* := i_{\tilde{E}}(E) \subset \tilde{E}$ .

**Proposition 5.7.** *For  $k \in \mathbb{Z}_{>0} \cup \{\infty\}$  the map*

$$\begin{aligned} \theta^k : \mathbf{G}(k) \cong \mathbf{G}(W^{(-k)}, E, W) &\hookrightarrow \mathbf{GO}(W, \tilde{E}, V) \cong \mathbf{GO}^1(\infty, 0) \\ U &\mapsto U \oplus (U^\perp \cap \text{span}(E_*)) \end{aligned}$$

is a linear embedding of linear ind-Grassmannians.

*Proof.* For any finite  $k$ , the embedding  $\theta^k$  is obtained as the direct limit  $\theta^k := \lim_{\rightarrow} \theta_m^k$  of the embeddings  $\theta_m^k$  defined in Subsection 3.1. For  $k = \infty$ , the embedding  $\theta^\infty : \mathbf{G}(\infty) \hookrightarrow \mathbf{GO}^1(\infty, 0)$  is defined as the direct limit  $\theta^\infty := \lim_{\rightarrow} \theta_{4l}^{2l}$ . □

### 5.3. Classification of linear embeddings.

**Theorem 5.8.** (Linear embeddings of linear ind-Grassmannians of the same type)

Let  $\varphi : \mathbf{X} \rightarrow \mathbf{Y}$  be a linear embedding of linear ind-Grassmannians, which does not factor through a projective ind-space.

- (i) If  $\varphi$  has the form  $\varphi : \mathbf{G}(m) \hookrightarrow \mathbf{G}(l)$  for  $l, m \in \mathbb{N} \cup \{\infty\}$ , then  $\varphi$  is a standard extension.
- (ii.1) If  $\varphi$  has the form  $\varphi : \mathbf{GO}(m, \infty) \hookrightarrow \mathbf{GO}(l, \infty)$  for  $m, l \in \mathbb{N} \cup \{\infty\}$ , then there are the following options:
  - $\varphi$  is a standard extension and  $m \leq l$  holds;
  - $\varphi$  is a combination of standard and isotropic extensions;
  - $\varphi$  factors through a standard ind-quadric.
- (ii.2) If  $\varphi$  has the form  $\varphi : \mathbf{GO}^\varepsilon(\infty, c) \hookrightarrow \mathbf{GO}^{\varepsilon'}(\infty, d)$  for  $c, d \in \mathbb{N}$ ,  $\varepsilon, \varepsilon' \in \{0, 1\}$  with the additional requirement  $c + \varepsilon \neq 2$  if  $d = 0$ , then there are the following options:
  - $\varphi$  is a standard extension and  $d \geq c$  holds;
  - $c \neq 0 \neq d$  and  $\varphi$  is a combination of standard and isotropic extensions;
  - $c > 0 = d$  and  $\varphi$  is a composition  $\theta^k \circ \sigma \circ \tau$  where  $\tau$  is a tautological embedding,  $\sigma$  is a standard extension, and  $k \geq 2$ ;
  - $c = 1$  and  $\varphi$  factors as  $\iota \circ \sigma \circ \delta^1$  or as  $\sigma_O \circ \delta_O^1$ , where  $\delta^1, \delta_O^1$  are defined in Proposition 5.6,  $\sigma, \sigma_O$  are standard extensions and  $\iota$  is an isotropic extension;
  - $c = 2$  and  $\varphi$  factors as  $\iota \circ \sigma \circ \delta^0$  or as  $\sigma_O \circ \delta_O^0$ , where  $\delta^0, \delta_O^0$  are defined in Proposition 5.6,  $\sigma, \sigma_O$  are standard extensions and  $\iota$  is an isotropic extension.
- (ii.3) If  $\varphi$  has the form  $\varphi : \mathbf{GO}^\varepsilon(\infty, c) \hookrightarrow \mathbf{GO}(l, \infty)$  for  $c \in \mathbb{N}$ ,  $l \in \mathbb{N} \cup \{\infty\}$ , then there are the following options:
  - $\varphi$  factors through a standard ind-quadric;
  - $c \neq 0$  and  $\varphi$  is a combination of standard and isotropic extensions;
  - $c = 1$  and  $\varphi$  factors as  $\iota \circ \sigma \circ \delta^1$ , or as  $\sigma_O \circ \delta_O^1$ , where  $\delta^1, \delta_O^1$  are defined in Proposition 5.6,  $\sigma, \sigma_O$  are standard extensions and  $\iota$  is an isotropic extension;
  - $c = 2$  and  $\varphi$  factors as  $\iota \circ \sigma \circ \delta^0$ , or as  $\sigma_O \circ \delta_O^0$ , where  $\delta^0, \delta_O^0$  are defined in Proposition 5.6,  $\sigma, \sigma_O$  are standard extensions and  $\iota$  is an isotropic extension.
- (ii.4) If  $\varphi$  has the form  $\varphi : \mathbf{GO}(m, \infty) \hookrightarrow \mathbf{GO}^\varepsilon(\infty, d)$  for  $m \in \mathbb{Z}_{\geq 1} \cup \{\infty\}$ ,  $d \in \mathbb{N}$  and  $\varepsilon \in \{0, 1\}$ , then  $\varphi$  is a combination of standard and isotropic extensions for  $d \neq 0$  and a composition  $\theta^k \circ \sigma \circ \tau$  of a tautological embedding  $\tau$ , a standard extension  $\sigma$  and one of the embeddings  $\theta^k$  for  $d = 0$ .
- (iii) If  $\varphi$  has the form  $\varphi : \mathbf{GS}(m, c) \hookrightarrow \mathbf{GS}(l, d)$  for  $m, l, c, d \in \mathbb{N} \cup \{\infty\}$ ,  $\{\infty\} \in \{m, c\} \cap \{l, d\}$ , then there are the following options:
  - $\varphi$  is a standard extension;
  - $\varphi$  is a combination of standard and isotropic extensions.

**Theorem 5.9.** (Linear embeddings of linear ind-Grassmannians of different types)

Let  $\varphi : \mathbf{X} \rightarrow \mathbf{Y}$  be a linear embedding of linear ind-Grassmannians, which does not factor through a projective ind-space.

- (i.1) If  $\varphi$  has the form  $\varphi : \mathbf{G}(m) \hookrightarrow \mathbf{GO}(l, \infty)$ , then  $\varphi$  is a composition

$$\mathbf{G}(m) \xrightarrow{\sigma} \mathbf{G}(l) \xrightarrow{\iota} \mathbf{GO}(l, \infty)$$

where  $\sigma$  is a standard extension and  $\iota$  is isotropic, or  $\varphi$  factors through a standard ind-quadric.

- (i.2) If  $\varphi$  has the form  $\varphi : \mathbf{G}(m) \hookrightarrow \mathbf{GO}^\varepsilon(\infty, l)$ , then  $\varphi$  is a composition

$$\mathbf{G}(m) \xrightarrow{\sigma} \mathbf{G}(k) \xrightarrow{\iota} \mathbf{GO}^\varepsilon(\infty, l)$$

where  $\sigma$  is a standard extension, and  $\iota$  is isotropic with  $k = l$  for  $l > 0$  and is the embedding  $\theta^k$  for  $l = 0$ .

(ii) If  $\varphi$  has the form  $\varphi : \mathbf{G}(m) \hookrightarrow \mathbf{GS}(l)$ , then  $\varphi$  is a composition

$$\mathbf{G}(m) \xrightarrow{\sigma} \mathbf{G}(l) \xrightarrow{\iota} \mathbf{GS}(l, \infty)$$

where  $\sigma$  is a standard extension and  $\iota$  is isotropic.

(iii.1) If  $\varphi$  has the form  $\varphi : \mathbf{GO}(m, \infty) \hookrightarrow \mathbf{G}(l)$ , then  $\varphi$  is a composition

$$\mathbf{GO}(m, \infty) \xrightarrow{\tau} \mathbf{G}(m) \xrightarrow{\sigma} \mathbf{G}(l)$$

where  $\tau$  is a tautological embedding and  $\sigma$  is a standard extension.

(iii.2) If  $\varphi$  has the form  $\varphi : \mathbf{GO}^\varepsilon(\infty, m) \hookrightarrow \mathbf{G}(l)$ , then  $\varphi$  is a composition

$$\mathbf{GO}^\varepsilon(\infty, m) \xrightarrow{\tau} \mathbf{G}(\infty) \xrightarrow{\sigma} \mathbf{G}(\infty)$$

where  $\tau$  is a tautological embedding and  $\sigma$  is a standard extension, or  $m = 2 - \varepsilon$  holds and the embedding  $\varphi$  is a composition

$$\mathbf{GO}^\varepsilon(\infty, 2 - \varepsilon) \xrightarrow{\delta^\varepsilon} \mathbf{G}(2) \xrightarrow{\sigma} \mathbf{G}(l)$$

where  $\delta^\varepsilon$  is given in Proposition 5.6 and  $\sigma$  is a standard extension.

(iv) If  $\varphi$  has the form  $\varphi : \mathbf{GS}(m, c) \hookrightarrow \mathbf{G}(l)$ , then  $\varphi$  factors as a composition

$$\mathbf{GS}(m, \infty) \xrightarrow{\tau} \mathbf{G}(m) \xrightarrow{\sigma} \mathbf{G}(l) ,$$

where  $\tau$  is a tautological embedding and  $\sigma$  is a standard extension.

(v.1) If  $\varphi$  has the form  $\varphi : \mathbf{GO}(m, \infty) \hookrightarrow \mathbf{GS}(l, d)$ , then  $\varphi$  is a mixed combination of standard and isotropic extensions.

(v.2) If  $\varphi$  has the form  $\varphi : \mathbf{GO}^\varepsilon(\infty, m) \hookrightarrow \mathbf{GS}(l, d)$ , then  $l = d = \infty$  and  $\varphi$  is a mixed combination of standard and isotropic extensions, or  $m = 2 - \varepsilon$  holds and the embedding  $\varphi$  is a composition

$$\mathbf{GO}^\varepsilon(\infty, 2 - \varepsilon) \xrightarrow{\delta_S^\varepsilon} \mathbf{GS}(2, \infty) \xrightarrow{\psi} \mathbf{GS}(l, d) ,$$

where  $\delta_S^\varepsilon$  is given in Proposition 5.6 and  $\psi$  is either a standard extension or a combination of standard and isotropic extensions.

(vi) If  $\varphi$  has the form  $\varphi : \mathbf{GS}(m, c) \hookrightarrow \mathbf{GO}(l, \infty)$ , then  $\varphi$  is a mixed combination of standard and isotropic extensions, or factors through a standard ind-quadric. If  $\varphi$  has the form  $\varphi : \mathbf{GS}(m, c) \hookrightarrow \mathbf{GO}^\varepsilon(\infty, d)$  with the additional requirement  $c \neq 0$  for  $d = 0$ , then  $\varphi = \iota \circ \sigma \circ \tau$  where  $\tau$  is a tautological embedding,  $\sigma$  is a standard extension, and  $\iota$  is an isotropic extension for  $d > 0$  and one of the embeddings  $\theta^k$  for  $d = 0$ .

*Proof of Theorems 5.8 and 5.9.* Let  $\varphi : \mathbf{X} \hookrightarrow \mathbf{Y}$  be obtained as the direct limit of a sequence of linear embeddings  $\varphi_k : X_k \hookrightarrow Y_k$  for appropriate standard exhaustions  $\mathbf{X} = \varinjlim X_k$  and  $\mathbf{Y} = \varinjlim Y_k$ . Thus we have a commutative diagram

$$(21) \quad \begin{array}{ccccccc} \dots & \xrightarrow{\psi_{k-1}} & X_k & \xrightarrow{\psi_k} & X_{k+1} & \xrightarrow{\psi_{k+1}} & \dots \\ & & \varphi_k \downarrow & & \varphi_{k+1} \downarrow & & \\ \dots & \xrightarrow{\tilde{\psi}_{k-1}} & Y_k & \xrightarrow{\tilde{\psi}_k} & Y_{k+1} & \xrightarrow{\tilde{\psi}_{k+1}} & \dots \end{array}$$

where  $\psi_k$  and  $\tilde{\psi}_k$  are standard extensions.

We start by addressing part (i) of Theorem 5.8. Here  $\mathbf{X} := \mathbf{G}(m) = \varinjlim G(m_k, V_{n_k})$  and  $\mathbf{Y} := \mathbf{G}(l) = \varinjlim G(l_k, \tilde{V}_{s_k})$ . Since  $\varphi$  does not factor through a projective ind-space, Theorem 2.4 implies that the embeddings  $\varphi_k$  exhausting  $\varphi$  are standard extensions. Furthermore, we reduce to the case where all embeddings in the diagram (21) are strict standard extensions, by omitting if necessary some embeddings  $\varphi_k$  and/or replacing each  $V_{n_k}$  by  $V_{n_k}^*$  and/or replacing each  $V_{n_k}$  by  $V_{n_k}^*$ .

To show that  $\varphi$  is a standard extension as in (19) we need to construct a global expression for  $\varphi$  from the local data of the above exhaustions. Recall first that every strict standard extension of ordinary Grassmannians  $G(m, V_n) \xrightarrow{\sigma} G(l, V_s)$  corresponds to an inclusion of vector spaces  $\nu : V_n \hookrightarrow V_s$ . In fact, the inclusion  $\nu$  is determined by  $\sigma$ . Indeed, the pullback  $\sigma^* \mathcal{S}_{G(l, V_s)}^*$  splits as a direct sum of  $\mathcal{S}_{G(m, V_n)}^*$  and a trivial bundle, and the inclusion  $\nu$  is dual to the surjection

$$V_s^* \cong H^0(G(l, V_s), \mathcal{S}_{G(l, V_s)}^*) \xrightarrow{\sim} H^0(G(m, V_n), \sigma^* \mathcal{S}_{G(l, V_s)}^*) \rightarrow H^0(G(m, V_n), \mathcal{S}_{G(m, V_n)}^*) \cong V_n^* .$$

By applying this to the strict standard extensions in the diagram (21), we obtain a commutative diagram of inclusions  $\nu_k, \mu_k, \tilde{\mu}_k$  of vector spaces

$$(22) \quad \begin{array}{ccccccc} \dots & \xrightarrow{\mu_{k-1}} & V_{n_k} & \xrightarrow{\mu_k} & V_{n_{k+1}} & \xrightarrow{\mu_{k+1}} & \dots \\ & & \nu_k \downarrow & & \nu_{k+1} \downarrow & & \\ \dots & \xrightarrow{\tilde{\mu}_{k-1}} & \tilde{V}_{s_k} & \xrightarrow{\tilde{\mu}_k} & \tilde{V}_{s_{k+1}} & \xrightarrow{\tilde{\mu}_{k+1}} & \dots \end{array} .$$

Now the direct limit  $\nu := \varinjlim \nu_k$  of the inclusions  $\nu_k : V_{n_k} \hookrightarrow \tilde{V}_{s_k}$  is an inclusion of countable-dimensional vector spaces

$$\nu : V := \varinjlim V_{n_k} \hookrightarrow \tilde{V} := \varinjlim \tilde{V}_{s_k} .$$

To complete the global expression of  $\varphi$  we need to find suitable subspaces and bases of  $V$  and  $\tilde{V}$ . For each  $k$ , the embedding  $\varphi_k$  is written as

$$\varphi_k(U) = \nu_k(U) \oplus W^{(k)} , \quad \text{where} \quad W^{(k)} := \bigcap_{U \in G(m_k, V_{n_k})} \varphi_k(U) .$$

Since each  $W^{(k)}$  is naturally a subspace of  $\tilde{V}$ , we can define the intersection

$$\underline{W}^{(k)} := \bigcap_{j \geq k} W^{(j)} .$$

Then  $\underline{W}^{(k-1)} \subset \underline{W}^{(k)}$  and the direct limit  $W' := \varinjlim \underline{W}^{(k)}$  is identified with a subspace of  $\tilde{V}$  satisfying

$$\nu(V) \cap W' = 0 \quad , \quad W' = \bigcap_{U \in \mathbf{X}} \varphi(U) .$$

Next, we fix a sequence  $U_{m_k} \in G(m_k, V_{n_k})$  such that  $\psi_k(U_{m_k}) = U_{m_{k+1}}$  for all  $k$ , and define subspaces  $W \subset V$  and  $\tilde{W} \subset \tilde{V}$  by setting  $W := \varinjlim U_{m_k}$  and  $\tilde{W} := \nu(W) \oplus W'$ . It remains to construct relevant bases of  $V$  and  $\tilde{V}$ . We start with a basis of  $E_W$  of  $W$  such that  $E_W \cap U_{m_k}$  is a basis of  $U_{m_k}$  for every  $k$ . Then we extend  $E_W$  to a basis  $E$  of  $V$  such that  $E \cap V_{n_k}$  is a basis of  $V_{n_k}$ . We fix a basis  $E'$  of  $W'$  such that  $E' \cap \underline{W}^{(k)}$  is a basis of  $\underline{W}^{(k)}$  for all  $k$ . Then  $\nu(E) \sqcup E'$  is a basis of  $\nu(V) \oplus W'$ , and  $\nu(E_W) \sqcup E'$  is a basis of  $\tilde{W}$ . We extend  $\nu(E) \sqcup E'$  to a basis  $\tilde{E}$  of  $\tilde{V}$ .

We have now constructed identifications  $\mathbf{X} = \mathbf{G}(W, E, V)$  and  $\mathbf{Y} = \mathbf{G}(\tilde{W}, \tilde{E}, \tilde{V})$  such that the embedding  $\varphi$  is written as

$$(23) \quad \begin{array}{ccc} \varphi : \mathbf{G}(m) = \mathbf{G}(W, E, V) & \hookrightarrow & \mathbf{G}(\tilde{W}, \tilde{E}, \tilde{V}) = \mathbf{G}(l) . \\ U & \mapsto & \nu(U) \oplus W' \end{array}$$

Hence  $\varphi$  is a standard extension. This completes the proof of part (i) of Theorem 5.8.

To address the linear embeddings between orthogonal ind-Grassmannians (parts (ii.1)-(ii.4) of Theorem 5.8), we first need to classify linear embeddings between ordinary ind-Grassmannians and orthogonal ind-Grassmannians (parts (i.1),(i.2),(iii.1),(iii.2) of Theorem 5.9).

For part (i.1) of Theorem 5.9, the embedding  $\varphi$  has the form

$$\varphi : \mathbf{G}(m) \cong \mathbf{X} \hookrightarrow \mathbf{Y} \cong \mathbf{GO}(l, \infty) .$$

By Theorem 2.5 each of the embeddings  $\varphi_k$  factors as a composition of a standard extension  $\sigma_k$  and an isotropic extension  $\iota_k$ :

$$\varphi_k : X_k = G(m_k, V_{n_k}) \xrightarrow{\sigma_k} G(l_k, \hat{V}_{r_k}) \xrightarrow{\iota_k} GO(l_k, \tilde{V}_{s_k}) = Y_k$$

for appropriate spaces  $\hat{V}_{r_k}$ . We take the canonical choice for  $\hat{V}_{r_k}$  given as the sum of the subspaces  $\varphi_k(U)$  for  $U \in X_k$ , which is an isotropic subspace of  $\tilde{V}_{s_k}$ .

Similarly to a standard extension of ordinary Grassmannians, the standard extensions  $\tilde{\psi}_k : GO(l_k, \tilde{V}_{s_k}) \hookrightarrow GO(l_{k+1}, \tilde{V}_{s_{k+1}})$  induce inclusions  $\tilde{\mu}_k : \tilde{V}_{s_k} \hookrightarrow \tilde{V}_{s_{k+1}}$ . We have  $\tilde{\mu}_k(\hat{V}_{r_k}) \subset \hat{V}_{r_{k+1}}$  due to the commutativity relation  $\varphi_{k+1} \circ \psi_k = \tilde{\psi}_k \circ \varphi_k$ . Therefore the restriction of  $\tilde{\psi}_k$  to  $G(l_k, \hat{V}_{r_k})$  defines a strict standard extension  $\eta_k : G(l_k, \hat{V}_{r_k}) \hookrightarrow G(l_k, \hat{V}_{r_{k+1}})$ . It follows that  $\varphi$  factors through the ind-Grassmannian  $\mathbf{Z} := \lim_{\rightarrow} G(l_k, \hat{V}_{r_k}) \cong \mathbf{G}(l)$  as

$$\mathbf{X} \xrightarrow{\sigma} \mathbf{Z} \xrightarrow{\iota} \mathbf{Y}$$

where  $\sigma := \lim_{\rightarrow} \sigma_k$ , and  $\iota$  is an isotropic extension. To complete the argument, we can apply part (i) of Theorem 5.8 to deduce that  $\sigma$  is a standard extension as required.

The proofs of the remaining statements of Theorems 5.8 and 5.9 follow the same general lines, and we just give the details in one least straightforward case: statement (iii.2) for  $\mathbf{X} \cong \mathbf{GO}^0(\infty, 2)$  and  $\mathbf{Y} \cong \mathbf{G}(l)$  with  $l \in \mathbb{Z}_{\geq 2} \cup \{\infty\}$ .

Here  $X_k \cong GO(n_k - 2, V_{2n_k})$  and  $Y_k \cong G(l_k, \tilde{V}_{r_k})$ . Since none of the embeddings  $\varphi_k$  factors through a projective space, by Theorem 3.3 there are two options for each  $\varphi_k$ : it factors as a composition

$$GO(n_k - 2, V_{2n_k}) \xrightarrow{\tau_k} G(n_k - 2, V_{2n_k}) \xrightarrow{\sigma_k^1} G(l_k, \tilde{V}_{r_k})$$

where  $\tau_k$  is the tautological embedding of  $X_k$  and  $\sigma_k^1$  is a standard extension, or  $\varphi_k$  factors as a composition

$$GO(n_k - 2, V_{2n_k}) \xrightarrow{\delta_{n_k}} G(2, S_{2n_k}) \xrightarrow{\sigma_k^2} G(l_k, \tilde{V}_{r_k})$$

where  $S_{2n_k} := GO(n_k, V_{2n_k})$ ,  $\delta_{n_k}$  is the linear embedding defined in (8), and  $\sigma_k^2$  is a standard extension. Clearly we can assume that one of the these two options for  $\varphi_k$  holds simultaneously for all  $k$ . We consider the two cases separately.

Suppose first that  $\varphi_k$  factors through  $\tau_k$  for all  $k$ . In this case we necessarily have  $l = \lim_{k \rightarrow \infty} l_k = \infty$ . For each  $k$  there is an isomorphism  $V_{2n_k} \cong V_{2n_k}^*$  provided by the respective symmetric bilinear form. This allows us to assume that the standard extension  $\sigma_k^1 : G(n_k - 2, V_{2n_k}) \hookrightarrow G(l_k, \tilde{V}_{r_k})$  is strict for all  $k$ , and so are the standard extensions  $\tilde{\psi}_k$ . Let  $\mu_k : V_{2n_k} \hookrightarrow V_{2n_{k+1}}$  be the inclusions associated to the standard extensions  $\psi_k$ . Then there are strict standard extensions  $\xi_k$  such that the following diagram is commutative for all  $k$ :

$$\begin{array}{ccccc} \varphi_{k+1} : & GO(n_{k+1} - 2, V_{2n_{k+1}}) & \xrightarrow{\tau_{k+1}} & G(n_{k+1} - 2, V_{2n_{k+1}}) & \xrightarrow{\sigma_{k+1}^1} & G(l_{k+1}, \tilde{V}_{r_{k+1}}) \\ & \psi_k \uparrow & & \xi_k \uparrow & & \tilde{\psi}_k \uparrow \\ \varphi_k : & GO(n_k - 2, V_{2n_k}) & \xrightarrow{\tau_k} & G(n_k - 2, V_{2n_k}) & \xrightarrow{\sigma_k^1} & G(l_k, \tilde{V}_{r_k}). \end{array}$$

Set  $V := \lim_{\rightarrow} V_{2n_k}$ , and let  $W^{(-2)} := \lim_{\rightarrow} U_{n_k-2}$  be the subspace of  $V$  defined by a chain  $U_{n_k-2} \in X_k$  such that  $\psi_k(U_{n_k-2}) = U_{n_{k+1}-2}$ . Let  $E \subset V$  be a isotropic basis containing a basis of each of the subspaces  $V_{2n_k}$ ,  $U_{n_k-2}$  for all  $k$ . Then the direct limit of the tautological embeddings  $\tau_k$  admits the following global expression:

$$\begin{array}{ccc} \tau_{\mathbf{X}} = \lim_{\rightarrow} \tau_k : \mathbf{X} = \mathbf{GO}(W^{-2}, E, V) & \hookrightarrow & \mathbf{G}(W, E, V) \cong \lim_{\rightarrow} G(n_k - 2, V_{2n_k}) = \mathbf{G}(\infty). \\ & & U \mapsto U \end{array}$$

Since each  $\sigma_k^1$  is a strict standard extension, the corresponding inclusion  $\nu_k : V_{2n_k} \hookrightarrow \tilde{V}_{r_k}$  is well defined, and we have an inclusion of direct limits  $\nu := \lim_{\rightarrow} \nu_k : V \hookrightarrow \tilde{V} := \lim_{\rightarrow} \tilde{V}_{r_k}$ . The argument used for part (i) of Theorem 5.8 implies that the direct limit  $\sigma^1 := \lim_{\rightarrow} \sigma_k^1 : \mathbf{G}(\infty) \rightarrow \mathbf{G}(\infty)$  is a standard extension of the form

$$\begin{array}{ccc} \sigma^1 = \lim_{\rightarrow} \sigma_k^1 : \mathbf{G}(W, E, V) & \hookrightarrow & \mathbf{G}(\tilde{W}, \tilde{E}, \tilde{V}) = \mathbf{Y} \\ & & U \mapsto U \oplus W' \end{array}$$

for some suitable subspace  $W' \subset \tilde{W}$  and basis  $E$ . We conclude that the initial embedding  $\varphi$  factors as  $\varphi = \sigma^1 \circ \tau_{\mathbf{X}}$  as asserted.

Next, suppose that  $\varphi_k$  factors through  $\delta_{n_k}$  for all  $k$ . We claim that there exist standard extensions  $\eta_k$  making the following diagram commutative for every  $k$ :

$$(24) \quad \begin{array}{ccccc} \varphi_{k+1} : GO(n_{k+1} - 2, V_{2n_{k+1}}) & \xrightarrow{\delta_{n_{k+1}}} & G(2, V_{2^{n_{k+1}-1}}) & \xrightarrow{\sigma_{k+1}^2} & G(l_{k+1}, \tilde{V}_{r_{k+1}}) \\ & \psi_k \uparrow & \eta_k \uparrow & & \tilde{\psi}_k \uparrow \\ \varphi_k : GO(n_k - 2, V_{2n_k}) & \xrightarrow{\delta_{n_k}} & G(2, V_{2^{n_k-1}}) & \xrightarrow{\sigma_k^2} & G(l_k, \tilde{V}_{r_k}). \end{array}$$

Note first that, as in the previous case, we can modify the exhaustions of  $\mathbf{X}$  and  $\mathbf{Y}$  so that the standard extensions  $\sigma_k^2$  and  $\tilde{\psi}_k$  are strict for all  $k$ . Each standard extension  $\psi_k : GO(n_k - 2, V_{2n_k}) \hookrightarrow GO(n_{k+1} - 2, V_{2n_{k+1}})$  has the form  $\psi_k(U) = \mu_k(U) \oplus U'$  for a unique  $U' \in GO(n_{k+1} - n_k, V_{2n_k})$ . Thus  $\psi_k$  determines a standard extension

$$\zeta_k : S_{2n_k} := GO(n_k, V_{2n_k}) \hookrightarrow GO(n_{k+1}, V_{2n_{k+1}}) =: S_{2n_{k+1}}, \quad \zeta_k(U) = \mu_k(U) \oplus U'.$$

Let  $\hat{\zeta}_k : V_{S_{2n_k}} \hookrightarrow V_{S_{2n_{k+1}}}$  be the inclusion corresponding to the embedding  $\zeta_k$ . Then  $\hat{\zeta}_k$  determines a strict standard extension

$$\eta_k : G(2, V_{S_{2n_k}}) \hookrightarrow G(2, V_{S_{2n_{k+1}}}), \quad \eta_k(U) := \hat{\zeta}_k(U).$$

Furthermore, we have  $\delta_{n_{k+1}} \circ \psi_k(U) = \hat{\zeta}_k(\delta_k(U))$ , which implies that the commutation relation  $\eta_k \circ \delta_{n_k} = \delta_{n_{k+1}} \circ \psi_k$  is satisfied for every  $k$ . Next, the construction from Case 2 of the proof of Theorem 3.3,(i), provides a linear embedding  $\pi_k : S_{n_k} \hookrightarrow \mathbb{P}(V_{r_k})$  such that the standard extension  $\sigma_k^2$  is induced by  $\pi_k$ . Let  $\hat{\pi}_k : V_{S_{n_k}} \hookrightarrow V_{r_k}$  be the inclusion induced by  $\pi_k$ . Then the relation  $\varphi_{k+1} \circ \psi_k = \tilde{\psi}_k \circ \varphi_k$  implies  $\hat{\pi}_{k+1} \circ \hat{\zeta}_k = \tilde{\mu}_k \circ \hat{\pi}_k$ , and hence  $\tilde{\psi}_k \circ \sigma_k^2 = \sigma_{k+1}^2 \circ \eta_k$ . This shows that the diagram (24) is commutative.

The existence of the standard extensions  $\eta_k$  implies that the embedding  $\varphi$  factors through the linear ind-Grassmannian  $\mathbf{G}(2) := \lim_{\rightarrow} G(2, V_{S_{2n_k}})$  as  $\varphi = \delta \circ \sigma^2$  where  $\delta := \lim_{\rightarrow} \delta_{n_k}$  and  $\sigma^2 := \lim_{\rightarrow} \sigma_k^2$ . Global expressions for  $\delta$  and  $\sigma^1$  are constructed respectively in Proposition 5.6 and in the proof of part (i) of Theorem 5.8. The classification of linear embeddings  $\mathbf{GO}^0(\infty, 2) \hookrightarrow \mathbf{G}(l)$  (with  $2 \leq l \leq \infty$ ) is complete.

All remaining cases in the proof of Theorems 5.8 and 5.9 are left to the reader.  $\square$

**Proposition 5.10.** *Let  $Y$  be a linear ind-Grassmannian not isomorphic to  $\mathbf{GO}^1(\infty, 0)$ . Then every linear embedding  $\mathbf{GO}^1(\infty, 0) \hookrightarrow Y$  factors through a projective space, or possibly through a standard ind-quadric in case  $Y$  isomorphic to  $\mathbf{GO}(l)$  or  $\mathbf{GO}(\infty, \infty)$ .*

*Proof.* The statement follows from Proposition 3.1.  $\square$

Theorems 5.8 and 5.9 yield the following.

**Corollary 5.11.** *Let  $\mathbf{X} \xrightarrow{\varphi} \mathbf{Y}$  be a proper linear embedding of linear ind-Grassmannians which does not factor through a projective ind-space or an ind-quadric. Then the following is a complete list of the possible choices for  $\mathbf{X}$  and  $\mathbf{Y}$ :*

- $\varphi : \mathbf{G}(m) \hookrightarrow \mathbf{G}(l)$  for  $2 \leq m \leq l \leq \infty$ ;
- $\varphi : \mathbf{GO}(m, \infty) \hookrightarrow \mathbf{GO}(l, \infty)$  for  $2 \leq m \leq l \leq \infty$ ;
- $\varphi : \mathbf{GO}(m, \infty) \hookrightarrow \mathbf{GO}^\varepsilon(\infty, l)$  for  $2 \leq m \leq l < \infty$  or  $l = 0$ ;
- $\varphi : \mathbf{GO}^\varepsilon(\infty, m) \hookrightarrow \mathbf{GO}(\infty, \infty)$  for  $0 < m < \infty$ ;
- $\varphi : \mathbf{GO}^\varepsilon(\infty, m) \hookrightarrow \mathbf{GO}^{\varepsilon'}(\infty, l)$  for  $0 < m \leq l < \infty$  or  $l = 0$ ;
- $\varphi : \mathbf{GS}(m, \infty) \hookrightarrow \mathbf{GS}(l, d)$  for  $2 \leq m \leq \min\{l, d\}$ ;
- $\varphi : \mathbf{GS}(\infty, m) \hookrightarrow \mathbf{GS}(\infty, l)$  for  $0 \leq m \leq l \leq \infty$ ;
- $\varphi : \mathbf{G}(m) \hookrightarrow \mathbf{GO}(l, \infty)$  for  $2 \leq m \leq l$ ;
- $\varphi : \mathbf{G}(m) \hookrightarrow \mathbf{GO}^\varepsilon(\infty, l)$  for  $2 \leq m \leq l < \infty$  or  $l = 0$ ;
- $\varphi : \mathbf{GO}(m, \infty) \hookrightarrow \mathbf{G}(l)$  for  $2 \leq m \leq l$ ;
- $\varphi : \mathbf{GO}^\varepsilon(\infty, m) \hookrightarrow \mathbf{G}(l)$  for  $0 < m \leq \infty$  and  $l = \infty$ , or for  $m = 2 - \varepsilon \leq l - \varepsilon$ ;
- $\varphi : \mathbf{GS}(m, \infty) \hookrightarrow \mathbf{G}(l)$  for  $2 \leq m \leq l \leq \infty$ ;
- $\varphi : \mathbf{GS}(\infty, m) \hookrightarrow \mathbf{G}(\infty)$  for  $0 \leq m \leq l = \infty$ ;
- $\varphi : \mathbf{GO}(m, \infty) \hookrightarrow \mathbf{GS}(l, d)$  for  $2 \leq m \leq \min\{l, d\} \leq \infty$ ;
- $\varphi : \mathbf{GO}^\varepsilon(\infty, m) \hookrightarrow \mathbf{GS}(l, d)$  for  $l = d = \infty$ , or for  $m = 2 - \varepsilon \leq \min\{l, d\} - \varepsilon$ ;
- $\varphi : \mathbf{GS}(m, c) \hookrightarrow \mathbf{GO}(l, \infty)$  for  $2 \leq m \leq l \leq \infty$  and  $0 \leq c \leq \infty$ ;
- $\varphi : \mathbf{GS}(m, c) \hookrightarrow \mathbf{GO}^\varepsilon(\infty, l)$  for  $l = 0$  or  $2 \leq m \leq l < \infty$  and  $0 \leq c \leq \infty$ .

It is well known [1] that every linear ind-Grassmannian  $\mathbf{X}$  admits a transitive action of an ind-group  $\mathbf{G}$  obtained as a direct limit  $\varinjlim G_{X_k}$  where  $X_k \xrightarrow{\sigma_k} X_{k+1}$  is an exhaustion of  $\mathbf{X}$  by standard extensions. Furthermore, the image of the corresponding homomorphism  $G_{X_k} \xrightarrow{f_k} G_{X_{k+1}}$  is an  $\mathcal{LS}$ -subgroup for each  $k$ .

**Corollary 5.12.** *Let  $\mathbf{X}$  and  $\mathbf{Y}$  be a linear ind-Grassmannian. If  $\mathbf{Y}$  is a spinor ind-Grassmannian we assume in addition that  $\mathbf{X} \not\cong \mathbf{GS}(\infty, 0)$  and  $\mathbf{X} \not\cong \mathbf{GO}^\varepsilon(\infty, 2 - \varepsilon)$  for  $\varepsilon = 0, 1$ . Every linear embedding  $\mathbf{X} \xrightarrow{\varphi} \mathbf{Y}$ , which does not factor through a projective ind-space or a standard ind-quadric, is equivariant with respect to a homomorphism of ind-groups  $\mathbf{G} \xrightarrow{f} \mathbf{H}$  where  $\mathbf{G}$  acts transitively on  $\mathbf{X}$  and  $\mathbf{H}$  acts transitively on  $\mathbf{Y}$ .*

*Proof.* The equivariance properties of linear embeddings of Grassmannians shown in Theorem 4.1, Proposition 4.4 and Corollary 4.5, imply that for linear ind-Grassmannians the claimed equivariance holds for tautological embeddings, standard extensions, isotropic extensions, as well as for the embeddings  $\delta^\varepsilon, \delta_0^\varepsilon, \delta_S^\varepsilon$  with  $\varepsilon \in \{0, 1\}$ . This implies the result for all linear embeddings of ind-Grassmannians, due to the classification in Theorems 5.8 and 5.9.  $\square$

## REFERENCES

- [1] I. Dimitrov, I. Penkov, *Ind-varieties of generalized flags as homogeneous spaces for classical ind-groups*, IMRN No. 55 (2004), 2935-2953.
- [2] E. Dynkin, *Maximal subgroups of classical group*, Trudy Mosk. Mat. Ob.) **1** (1952), 39-166.
- [3] E. Dynkin, *Semisimple subalgebras of semisimple Lie algebras*, Mat. Sbornik (N.S.), **30**, (72) (1952), 349-462; English translation in Amer. Math. Soc. Transl., Series 2, Vol. 6 (1957), 111-244.
- [4] J.E. Humphreys, *Linear algebraic groups*, Springer-Verlag Berlin-Heidelberg-New York, 1975.
- [5] F. Knop, H. Kraft, and T. Vust, *The Picard group of a G-variety*, Algebraische Transformationsgruppen und Invariantentheorie, DMV Sem. **13**, 77–87, Birkhäuser, Basel, 1989.
- [6] J.M. Landsberg, *Tensors: Geometry and Applications*, Grad. Stud. in Math., Vol. **128**, AMS 2012.
- [7] J.M. Landsberg, L. Manivel, *On the projective geometry of rational homogeneous varieties*, Comment. Math. Helv. **78** (2003), 65-100.
- [8] A. Onishchik, *Inclusion relations between transitive compact transformation groups*, Tr. Mosk. Mat. Obs., **11** (1962), 199-242. In Russian; English Transl.: Amer. Math. Soc. Transl. **50** (1966), 5–58.
- [9] I. Penkov, A.S. Tikhomirov, *Linear ind-Grassmannians*, Pure and Applied Mathematics Quarterly **10** (2014), 289-323.
- [10] M. Sato, *The KP hierarchy and infinite-dimensional Grassmann manifolds*, in Theta functions — Bowdoin 1987, Part 1 (Brunswick, ME, 1987), Proc. Sympos. Pure Math. 49, Amer. Math. Soc., Providence, RI, 1989, 51–66.
- [11] J. Tits, *Sur les analogues algébriques des groupes semi-simples complexes*, in Colloque d’algèbre supérieure, Bruxelles, 19–22 décembre 1956, Centre Belge de Recherches Mathématiques, Établissements Ceuterick, Louvain (1957), 261–289.

Ivan Penkov  
 Constructor University, 28759 Bremen, Germany  
 E-mail address: ipenkov@constructor.university

Valdemar Tsanov  
 Institute of Mathematics and Informatics, Bulgarian Academy of Sciences,  
 Bulgaria, Sofia 1113, Acad. G. Bonchev Str., Bl. 8  
 and Constructor University, 28759 Bremen, Germany. E-mail address: valdemar.tsanov@math.bas.bg