

Foundations of Mathematical Physics

Homework 7

Due on Nov. 1, 2023, before the tutorial.

Problem 1 [5 points]: Orthonormal basis

Prove that an orthonormal sequence $(\varphi_j)_j$ in a Hilbert space is an orthonormal basis if and only if

$$\langle \varphi_j, \psi \rangle = 0 \quad \text{for all } j \in \mathbb{N} \quad \Rightarrow \quad \psi = 0.$$

Problem 2 [7 points]: Operator norm

Prove the following lemma that was stated in class: Let $\mathcal{L}(X, Y)$ be the set of bounded linear operators from $X \rightarrow Y$. Then $\mathcal{L}(X, Y)$ with the norm

$$\|L\|_{\mathcal{L}(X, Y)} := \sup_{\|x\|_X=1} \|Lx\|_Y$$

is a normed space. Furthermore, if Y is a Banach space, then so is $\mathcal{L}(X, Y)$.

Problem 3 [8 points]: Fourier transform

Let $f \in C_c^\infty(\mathbb{R}^d)$, and let $0 < \alpha < d$. Prove that then

$$c_\alpha \mathcal{F}^{-1}(|k|^{-\alpha} \widehat{f}(k))(x) = c_{d-\alpha} \int |x-y|^{\alpha-d} f(y) \, dy$$

for some constant c_α , and determine c_α explicitly. *Hint:* $\int_0^\infty e^{-\pi k^2 \lambda} \lambda^{\alpha/2-1} \, d\lambda = ?$. *You might want to look up or recall the definition of the gamma function.*

Note: In this sense we can give a meaning to the Fourier transform of $|x|^{\alpha-d}$.