

Last time: For $f \in C^{(n)}$, we write $f(x) = \sum_{k=0}^n \frac{f^{(k)}(c)}{k!} (x-c)^k + R_{n,c}(x)$.
 $\qquad\qquad\qquad := P_{n,c}(x) \qquad\qquad\qquad o((x-c)^n)$

Note: Taylor polynomials are unique: Let f be n times differentiable, and $c \in I$. If Q is a polynomial of degree $\leq n$ with $f(x) - Q(x) = o((x-c)^n)$ as $x \rightarrow c$, then $P_{n,c}(x) = Q(x)$.

\Rightarrow We can use any method to find a polynomial Q of degree $\leq n$ with

$f(x) = Q(x) + o((x-c)^n)$; then $Q(x)$ is always the Taylor polynomial.

(Useful, e.g., for products or compositions of fct.s) (\rightarrow see HW 1, Problem 3)

So far we have not made the remainder $R_{n,0}(x)$ very explicit. But using the fundamental thm. of calculus we can write:

$$\begin{aligned} f(x) - f(c) &= \underbrace{\int_c^x f'(t) dt}_{\text{int. by parts with } (t-x) \text{ as antiderivative of 1 (w.r.t. t)}} \quad \text{int. by parts with } (t-x) \text{ as antiderivative of 1 (w.r.t. t)} \\ &= \int_c^x 1 \cdot f'(t) dt = \underbrace{(t-x)f'(t) \Big|_c^x}_{= (x-c)f'(c)} - \int_c^x (t-x)f''(t) dt \\ &\qquad\qquad\qquad = (x-c)f'(c) \end{aligned}$$

$$\Rightarrow f(x) = f(c) + f'(c)(x-c) + \int_c^x (x-t)f''(t) dt$$

[check the next order or do a proof by induction]

Repeating this yields Taylor's thm.:

(let $f \in C^{(n+1)}(I)$, $I = (a,b)$, $c \in I$. If $x \in I$, then

$$R_{n,c}(x) = \int_c^x \frac{(x-t)^n}{n!} f^{(n+1)}(t) dt \quad (\text{integral form of remainder}).$$

This can be used for rigorous remainder estimates, e.g.:

- If additionally there is a $C > 0$ s.t. $|f^{(n+1)}(x)| \leq C \quad \forall x \in I$, then

$$|R_{n+1}(x)| \leq C \frac{|x-c|^{n+1}}{(n+1)!} = C \frac{|x-c|}{n+1} \cdot \frac{|x-c|}{n} \cdots \frac{|x-c|}{1} \xrightarrow{n \rightarrow \infty} 0 \quad \text{for any given } x, c.$$

$$\text{So in this case } f(x) = \lim_{n \rightarrow \infty} \sum_{k=0}^n \frac{f^{(k)}(c)}{k!} (x-c)^k \quad \text{for all } x \in I$$

Note: There are also the Cauchy and Lagrange forms of the remainder.

(see HW1 Problem 2 for the Lagrange form)

1.2 Sequences of Functions


Let $(f_n(x))_{n \in \mathbb{N}}$ be a sequence of functions $f_n: D \rightarrow \mathbb{R}$, $D \subset \mathbb{R}$.

Things we want to know (here $f(x) := \lim_{n \rightarrow \infty} f_n(x)$ for all $x \in D$):

- If all f_n are continuous, is f also continuous?
- Is $\lim_{n \rightarrow \infty} \int f_n(x) dx = \int \lim_{n \rightarrow \infty} f_n(x) dx = \int f(x) dx$?
- Is $\lim_{n \rightarrow \infty} f'_n(x) = f'(x)$?


Examples:

- $f_n(x) = x^n$, $x \in [0, 1]$



Each f_n is continuous, but $\lim_{n \rightarrow \infty} f_n(x) = \begin{cases} 0 & \text{for } x \in [0, 1) \\ 1 & \text{for } x = 1 \end{cases}$ is not continuous.

$$\cdot f_n(x) = \begin{cases} \frac{n^2}{2}x & ; 0 \leq x \leq \frac{1}{n} \\ 2n - n^2x & ; \frac{1}{n} \leq x \leq \frac{2}{n} \\ 0 & ; \frac{2}{n} \leq x \leq 1 \end{cases}$$



[check the computation]

Here, $\int_0^2 f_n(x) dx = \dots = 1$ for all $n \in \mathbb{N}$, but $\lim_{n \rightarrow \infty} f_n(x) = 0$ for all $x \in [0, 2]$.

So $\lim_{n \rightarrow \infty} \int_0^2 f_n(x) dx \neq \int_0^2 f(x) dx$.

$$\cdot f_n(x) = \frac{1}{n} \sin(nx).$$

Here, $f_n(x) \xrightarrow{n \rightarrow \infty} 0 = f(x)$, but $f'_n(x) = \cos(nx)$ does not have a limit $\forall x$ as $n \rightarrow \infty$.

Key concept to achieve nicer convergence results: uniform convergence

Above, we only used the notion of pointwise convergence:

$(f_n)_n$ converges to $f: D \rightarrow \mathbb{R}$ pointwise: $\Leftrightarrow f_n(x) \xrightarrow{n \rightarrow \infty} f(x)$ for all $x \in D$.

In other words:

For all $x \in D$ we have: $\forall \varepsilon > 0 \exists N \in \mathbb{N}$ s.t. $\forall n \geq N: |f_n(x) - f(x)| < \varepsilon$

Here, x is chosen first, and ε and N may depend on the choice of x .

Next we require ε and N that work for all x at the same time:


Definition:

(f_n) with $f_n: D \rightarrow \mathbb{R}$, $D \subset \mathbb{R}$ converges uniformly to $f: D \rightarrow \mathbb{R}$ if

$\forall \varepsilon > 0 \exists N \in \mathbb{N} \text{ s.t. } \forall n \geq N \text{ and } \forall x \in D: |f_n(x) - f(x)| < \varepsilon.$

here, the N large enough is independent of x

Graphically:



"all f_n 's are in any ε -tube around f (for n large enough)"

For the examples $f_n(x) = x^n$, $x \in [0, 1]$, and $f_n(x) = \begin{cases} n^2 x, & 0 \leq x \leq \frac{1}{n} \\ 2n - n^2 x, & \frac{1}{n} \leq x \leq \frac{2}{n} \\ 0, & \frac{2}{n} \leq x \leq 1 \end{cases}$
we do not have uniform convergence!