

Week 2: Elementary Analytical Geometry

1. ☐ MULTI ☐ Single

What is the angle (in radian, i.e., where 360° corresponds to 2π) between the vectors

$$\begin{bmatrix} 1 \\ 2 \\ -1 \end{bmatrix}$$

and

$$\begin{bmatrix} 3 \\ 7 \\ 17 \end{bmatrix}?$$

- (a) $\frac{\pi}{2}$
 (b) $\frac{\pi}{4}$
 (c) π
 (d) 0

2. ☐ MULTI ☐ Single

If $\vec{u}$ and $\vec{v}$ are perpendicular unit vectors, then

- (a) $|\vec{u} - \vec{v}| = 1$
 (b) $|\vec{u} - \vec{v}| = \sqrt{2}$
 (c) $|\vec{u} - \vec{v}| = 0$
 (d) $|\vec{u} - \vec{v}|$ cannot be computed without further information on $\vec{u}$ and $\vec{v}$

3. ☐ MULTI ☐ Single

How long is the vector $(1, 1, \dots, 1)$ in 16 dimensions?

- (a) Length = 1
 (b) Length = 4
 (c) Length = 16
 (d) Length = 32

4. ☐ MULTI ☐ Single

Which of the following formulas is not true (for $\vec{u}, \vec{v} \in \mathbb{R}^n$)?

- (a) $|\vec{u} \times \vec{v}| \leq |\vec{u}||\vec{v}|$
 (b) $\cos(\theta) = \frac{\vec{u} \cdot \vec{v}}{|\vec{u}|^2 |\vec{v}|^2}$
 (c) $|\vec{u} - \vec{v}|^2 = |\vec{u}|^2 - 2|\vec{u}||\vec{v}| \cos(\theta) + |\vec{v}|^2$
 (d) $|\vec{u} + \vec{v}| \leq |\vec{u}| + |\vec{v}|$

5. ☐ MULTI ☐ Single

Let x, y, z be such that $x + y + z = 0$. Define $u = (x, y, z)$ and $v = (z, x, y)$. What is the value of $\frac{u \cdot v}{|u||v|}$?

- (a) $\frac{-x^2 + yz}{x^2 + y^2 + (x - y)^2}$

- (b) $-\frac{1}{2}$
 (c) 1
 (d) 0

6. ☐ MULTI ☐ Single

A line is given by $\vec{r} = \lambda\vec{a} + \vec{b}$, with $\vec{a} = (1, -1, 4)$ and $\vec{b} = (4, 5, 6)$, while the equation of a plane is given by $-2x + 2y + z = 17$. What are the coordinates of the point P where the line and plane intersect?

- (a) The line and the plane intersect infinitely many times
 (b) $P = (3, 3, 17)$
 (c) The line and the plane do not intersect
 (d) $P = (-1, 4, 7)$

7. ☐ MULTI ☐ Single

What is the equation of the hyperplane, given by $\begin{bmatrix} t \\ x \\ y \\ z \end{bmatrix} = \vec{p}_0 + \alpha\vec{a} + \beta\vec{b} + \gamma\vec{c}$ with

$$\vec{p}_0 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \vec{a} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \vec{b} = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}, \vec{c} = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \end{bmatrix}, \alpha, \beta, \gamma \in \mathbb{R}$$

- (a) $t + x - y - z - 1 = 0$
 (b) $t + x - y + z - 1 = 0$
 (c) $-t - x - y - z + 1 = 0$
 (d) $-t - x - y + z + 1 = 0$

8. ☐ MULTI ☐ Single

Find the cross product $\vec{u} \times \vec{v}$ of $\vec{u} = \langle 3, 2, -1 \rangle$, $\vec{v} = \langle 1, 1, 0 \rangle$

- (a) $\langle -1, -1, 5 \rangle$
 (b) $\langle -6, -4, 2 \rangle$
 (c) $\langle 1, -1, 1 \rangle$
 (d) $\langle 6, -4, 2 \rangle$

9. ☐ MULTI ☐ Single

Find the unit vector along the direction of the cross product $\vec{u} \times \vec{v}$ of $\vec{u} = \langle 7, -1, 3 \rangle$, $\vec{v} = \langle 2, 0, -2 \rangle$.

- (a) $\frac{1}{\sqrt{108}} \langle -2, -10, 2 \rangle$
 (b) $\frac{1}{408} \langle 2, 20, 2 \rangle$
 (c) $\frac{1}{\sqrt{408}} \langle 2, 20, 2 \rangle$
 (d) $\frac{1}{108} \langle -2, -10, 2 \rangle$

10. MULTI Single

$$\text{Let } \epsilon_{ijk} = \begin{cases} 1 & \text{if } (i\ j\ k) = (1\ 2\ 3), (2\ 3\ 1), \text{ or } (3\ 1\ 2) \\ -1 & \text{if } (i\ j\ k) = (1\ 3\ 2), (3\ 2\ 1), \text{ or } (2\ 1\ 3) \\ 0 & \text{else} \end{cases}$$

Consider $\vec{u} = \langle u_1, u_2, u_3 \rangle$ and $\vec{v} = \langle v_1, v_2, v_3 \rangle$. Which of the following is equivalent to the k th component of $\vec{u} \times \vec{v}$

$$(a) \ [\vec{u} \times \vec{v}]_k = \sum_{i,j=1}^3 \epsilon_{ijk} u_i v_j$$

$$(b) \ [\vec{u} \times \vec{v}]_k = \sum_{i,j=1}^3 \epsilon_{ijk} (u_i v_j + v_i u_j)$$

$$(c) \ [\vec{u} \times \vec{v}]_k = \sum_{i,j=1}^3 \epsilon_{ijk} (u_i v_j - v_i u_j)$$

$$(d) \ [\vec{u} \times \vec{v}]_k = \sum_{i,j=1}^3 \epsilon_{ijk} v_i u_j$$

Total of marks: 10