

Elements of Calculus

Homework 7 (covering Weeks 13 and 14)

This homework cannot be submitted for grading, and solutions will be provided.

Problem 1 [4 points]

Find the general solution to the linear homogeneous ODE

$$y'' + y' - 2y = 0.$$

Then give the solution for the initial condition $y(0) = 2$ and $y'(0) = 5$. What is the behavior of the solution as $x \rightarrow \infty$? Also, find one particular solution to the linear inhomogeneous ODE

$$y'' + y' - 2y = e^{-x}.$$

Finally, provide the general solution (i.e., involving two constants) to this inhomogeneous ODE.

Problem 2 [6 points]

We consider the harmonic oscillator in a very general setting, i.e., with friction and a driving force. The corresponding ODE is

$$m \frac{d^2x}{dt^2} + r \frac{dx}{dt} + kx = f(t),$$

where $m, r, k > 0$, and $f(t)$ is an external driving force. We are looking for a solution $x(t)$ of this equation.

(a) Let us consider the homogeneous case first, i.e., $f = 0$.

- (1) Find the general real solution to the homogeneous equation. (You will need to consider different cases depending on m, r, k .)
- (2) How do the solutions behave for large t ?
- (3) What is the solution of the harmonic oscillator without friction, i.e., when $r = 0$?

(b) Now consider the inhomogeneous case with periodic force.

- (1) Find one particular complex solution to the inhomogeneous equation with $f(t) = e^{i\omega t}$.
- (2) By using Euler's formula, use part (1) to find one particular real solution to the inhomogeneous equation with $f(t) = \sin(\omega t)$.
- (3) For $f(t) = \sin(\omega t)$ and $r^2 < 2km$, which driving frequency ω makes the amplitude of the solution maximal? This is the so-called resonance frequency. What happens to the amplitude at resonance if the friction goes to zero, i.e., $r \rightarrow 0$?

Problem 3 [4 points]

Compute the Fourier transform of the “saw-tooth function” $f(x) = x$ on $[-\pi, \pi)$, periodically extended outside its fundamental domain.

Problem 4 [6 points]

- (a) Compute the Fourier series of the function $f(x) = (x - \pi)^2$ for $x \in [0, 2\pi]$ (Example B from Class Session 27).
- (b) Finish the proof that we started in class (Session 28) of the fact that the bump function from Example A (Session 27) is mean-square convergent by showing that

$$\sum_{k=-\infty}^{\infty} |\hat{f}_k|^2 = \frac{a}{2\pi},$$

where \hat{f}_k are the Fourier coefficients of the bump function.